



CITY OF CARMEL-BY-THE-SEA CLIMATE COMMITTEE

Contact: 831.620.2000 www.ci.carmel.ca.us

All meetings are held in the City Council Chambers
East Side of Monte Verde Street
Between Ocean and 7th Avenues

SPECIAL MEETING **Wednesday, August 18, 2021**

3:30 PM

Governor Newsom's Executive Order N-29-20 has allowed local legislative bodies to hold public meetings via teleconference and to make public meetings accessible telephonically or otherwise electronically to all members of the public seeking to observe and to address the local legislative body. Also, see the Order by the Monterey County Public Health Officer issued March 17, 2020. The health and well-being of our residents is the top priority for the City of Carmel-by-the-Sea. To that end, this meeting will be held via teleconference and web-streamed on the City's website ONLY.

To attend via Zoom <https://ci-carmel-ca-us.zoom.us/j/93340805428? Meeting ID 93340805428, Passcode 669209>; or to attend via telephone dial 1-312-626-6799. The public can also email comments to amartelet@ci.carmel.ca.us. Comments must be received 2 hours before the meeting in order to be provided to the committee. Comments received after that time and up to the beginning of the meeting will be added to the agenda and made part of the record.

Shortly after roll call, the Board/Commission will leave the Council Chambers for an on-site tour of inspection of the properties listed on the agenda. Prior to the beginning of the tour, the Board/Commission may eliminate one or more property. The public is welcome to join the tour. The Board/Commission will return to the Council Chambers after completing the tour to begin the meeting no earlier than the time noted below.

CALL TO ORDER

PUBLIC APPEARANCES

Members of the public are entitled to speak on matters of municipal concern not on the agenda during Public Appearances. Each person's comments shall be limited to 3 minutes, or as otherwise established by the Chair. Matters not appearing on the agenda will not receive action at this meeting and may be referred to staff. Persons are not required to provide their names, and it is helpful for speakers to state their names so they may be identified in the minutes of the meeting.

ANNOUNCEMENTS

ORDERS OF BUSINESS

Orders of Business are agenda items that require Committee discussion, debate, direction to staff, and/or action.

1. Review the IPCC Working Group 1 Summary for Policymakers
2. Update on Climate Action Plan Tasks and Next Steps
3. Review, provide comments, and/or approve the Summary Sheet for the Monterey Regional Waste Management District Presentation
4. Review and Update the List of Partners to Engage and Discuss the Public Outreach Approach for the Climate Action and Adaptation Strategy Development Phase

FUTURE AGENDA ITEMS

ADJOURNMENT

This agenda was posted at City Hall, Monte Verde Street between Ocean Avenue and 7th Avenue, outside the Park Branch Library, NE corner of Mission Street and 6th Avenue, the Carmel-by-the-Sea Post Office, 5th Avenue between Dolores Street and San Carlos Street, and the City's webpage <http://www.ci.carmel.ca.us> in accordance with applicable legal requirements.

SUPPLEMENTAL MATERIAL RECEIVED AFTER THE POSTING OF THE AGENDA

Any supplemental writings or documents distributed to a majority of the Climate Committee regarding any item on this agenda, received after the posting of the agenda will be available at the Public Works Department located on the east side of Junipero Street between Fourth and Fifth Avenues during normal business hours.

SPECIAL NOTICES TO PUBLIC

In compliance with the Americans with Disabilities Act, if you need special assistance to participate in this meeting, please contact the City Clerk's Office at 831-620-2000 at least 48 hours prior to the meeting to ensure that reasonable arrangements can be made to provide accessibility to the meeting (28CFR 35.102-35.104 ADA Title II).



CITY OF CARMEL-BY-THE-SEA

Climate Committee

Staff Report

August 18, 2021
ORDERS OF BUSINESS

TO: Climate Committee Members

SUBMITTED BY: Agnes Martelet, Environmental Compliance Manager

SUBJECT: Review the IPCC Working Group 1 Summary for Policymakers

RECOMMENDATION:

Review the IPCC Working Group 1 Summary for Policymakers

BACKGROUND/SUMMARY:

The Intergovernmental Panel on Climate Change (IPCC) released their Working Group I Summary for Policymakers on August 7, 2021. The Summary for policymakers presents key findings on the physical science basis of climate change. This is part of the IPCC's Sixth Assessment Report.

The report provides a science-based, high-level summary of the current state of the climate, how it is changing, human influence, and possible future paths for climate change. The report is included in Attachment 1.

The Climate Committee should review this report for information relevant to the City's climate action and adaptation planning efforts.

FISCAL IMPACT:


ATTACHMENTS:

Attachment 1: IPCC Working Group I Summary for Policymakers

Climate Change 2021

The Physical Science Basis

Summary for Policymakers



Summary for Policymakers

Attachment 1

Drafting Authors:

Richard P. Allan (United Kingdom), Paola A. Arias (Colombia), Sophie Berger (France/Belgium), Josep G. Canadell (Australia), Christophe Cassou (France), Deliang Chen (Sweden), Annalisa Cherchi (Italy), Sarah L. Connors (France/United Kingdom), Erika Coppola (Italy), Faye Abigail Cruz (Philippines), Aïda Diongue-Niang (Senegal), Francisco J. Doblas-Reyes (Spain), Hervé Douville (France), Fatima Driouech (Morocco), Tamsin L. Edwards (United Kingdom), François Engelbrecht (South Africa), Veronika Eyring (Germany), Erich Fischer (Switzerland), Gregory M. Flato (Canada), Piers Forster (United Kingdom), Baylor Fox-Kemper (United States of America), Jan S. Fuglestvedt (Norway), John C. Fyfe (Canada), Nathan P. Gillett (Canada), Melissa I. Gomis (France/Switzerland), Sergey K. Gulev (Russian Federation), José Manuel Gutiérrez (Spain), Rafiq Hamdi (Belgium), Jordan Harold (United Kingdom), Mathias Hauser (Switzerland), Ed Hawkins (United Kingdom), Helene T. Hewitt (United Kingdom), Tom Gabriel Johansen (Norway), Christopher Jones (United Kingdom), Richard G. Jones (United Kingdom), Darrell S. Kaufman (United States of America), Zbigniew Klimont (Austria/Poland), Robert E. Kopp (United States of America), Charles Koven (United States of America), Gerhard Krinner (France/Germany, France), June-Yi Lee (Republic of Korea), Irene Lorenzoni (United Kingdom/Italy), Jochem Marotzke (Germany), Valérie Masson-Delmotte (France), Thomas K. Maycock (United States of America), Malte Meinshausen (Australia/Germany), Pedro M.S. Monteiro (South Africa), Angela Morelli (Norway/Italy), Vaishali Naik (United States of America), Dirk Notz (Germany), Friederike Otto (United Kingdom/Germany), Matthew D. Palmer (United Kingdom), Izidine Pinto (South Africa/Mozambique), Anna Pirani (Italy), Gian-Kasper Plattner (Switzerland), Krishnan Raghavan (India), Roshanka Ranasinghe (The Netherlands/Sri Lanka, Australia), Joeri Rogelj (United Kingdom/Belgium), Maisa Rojas (Chile), Alex C. Ruane (United States of America), Jean-Baptiste Sallée (France), Bjørn H. Samset (Norway), Sonia I. Seneviratne (Switzerland), Jana Sillmann (Norway/Germany), Anna A. Sörensson (Argentina), Tannecia S. Stephenson (Jamaica), Trude Storelvmo (Norway), Sophie Szopa (France), Peter W. Thorne (Ireland/United Kingdom), Blair Trewin (Australia), Robert Vautard (France), Carolina Vera (Argentina), Nouredine Yassaa (Algeria), Sönke Zaehle (Germany), Panmao Zhai (China), Xuebin Zhang (Canada), Kirsten Zickfeld (Canada/Germany)

Contributing Authors:

Krishna M. AchutaRao (India), Bhupesh Adhikary (Nepal), Edvin Aldrian (Indonesia), Kyle Armour (United States of America), Govindasamy Bala (India/United States of America), Rondrotiana Barimalala (South Africa/Madagascar), Nicolas Bellouin (United Kingdom/France), William Collins (United Kingdom), William D. Collins (United States of America), Susanna Corti (Italy), Peter M. Cox (United Kingdom), Frank J. Dentener (EU/The Netherlands), Claudine Dereczynski (Brazil), Alejandro Di Luca (Australia, Canada/Argentina), Alessandro Dosio (Italy), Leah Goldfarb (France/United States of America), Irina V. Gorodetskaya (Portugal/Belgium, Russian Federation), Pandora Hope (Australia), Mark Howden (Australia), Akm Saiful Islam (Bangladesh), Yu Kosaka (Japan), James Kossin (United States of America), Svitlana Krakovska (Ukraine), Chao Li (China), Jian Li (China), Thorsten Mauritsen (Germany/Denmark), Sebastian Milinski (Germany), Seung-Ki Min (Republic of Korea), Thanh Ngo Duc (Vietnam), Andy Reisinger (New Zealand), Lucas Ruiz (Argentina), Shubha Sathyendranath (United Kingdom/Canada, Overseas Citizen of India), Aimée B. A. Slangen (The Netherlands), Chris Smith (United Kingdom), Izuru Takayabu (Japan), Muhammad Irfan Tariq (Pakistan), Anne-Marie Treguier (France), Bart van den Hurk (The Netherlands), Karina von Schuckmann (France/Germany), Cunde Xiao (China)

Date of Document: 7 August 2021 17:00 CEST

This Summary for Policymakers should be cited as:

IPCC, 2021: Summary for Policymakers. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press. In Press. Attachment 1

This document is subject to final copy-editing.

Table of Contents

Introduction	4	Attachment 1
A. The Current State of the Climate	5	
B. Possible Climate Futures	15	
C. Climate Information for Risk Assessment and Regional Adaptation.....	31	
D. Limiting Future Climate Change	36	

Introduction

Attachment 1

This Summary for Policymakers (SPM) presents key findings of the Working Group I (WGI) contribution to the IPCC's Sixth Assessment Report (AR6)¹ on the physical science basis of climate change. The report builds upon the 2013 Working Group I contribution to the IPCC's Fifth Assessment Report (AR5) and the 2018–2019 IPCC Special Reports² of the AR6 cycle and incorporates subsequent new evidence from climate science³.

This SPM provides a high-level summary of the understanding of the current state of the climate, including how it is changing and the role of human influence, the state of knowledge about possible climate futures, climate information relevant to regions and sectors, and limiting human-induced climate change.

Based on scientific understanding, key findings can be formulated as statements of fact or associated with an assessed level of confidence indicated using the IPCC calibrated language⁴.

The scientific basis for each key finding is found in chapter sections of the main Report, and in the integrated synthesis presented in the Technical Summary (hereafter TS), and is indicated in curly brackets. The AR6 WGI Interactive Atlas facilitates exploration of these key synthesis findings, and supporting climate change information, across the WGI reference regions⁵.

¹ Decision IPCC/XLVI-2.

² The three Special reports are: Global warming of 1.5°C: an IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty (SR1.5); Climate Change and Land: an IPCC Special Report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems (SRCCL); IPCC Special Report on the Ocean and Cryosphere in a Changing Climate (SROCC).

³ The assessment covers scientific literature accepted for publication by 31 January 2021.

⁴ Each finding is grounded in an evaluation of underlying evidence and agreement. A level of confidence is expressed using five qualifiers: very low, low, medium, high and very high, and typeset in italics, for example, *medium confidence*. The following terms have been used to indicate the assessed likelihood of an outcome or a result: *virtually certain* 99–100% probability, *very likely* 90–100%, *likely* 66–100%, *about as likely as not* 33–66%, *unlikely* 0–33%, *very unlikely* 0–10%, *exceptionally unlikely* 0–1%. Additional terms (*extremely likely* 95–100%, *more likely than not* >50–100%, and *extremely unlikely* 0–5%) may also be used when appropriate. Assessed likelihood is typeset in italics, for example, *very likely*. This is consistent with AR5. In this Report, unless stated otherwise, square brackets [x to y] are used to provide the assessed *very likely* range, or 90% interval.

⁵ The Interactive Atlas is available at <https://interactive-atlas.ipcc.ch>

A. The Current State of the Climate

Attachment 1

Since AR5, improvements in observationally based estimates and information from paleoclimate archives provide a comprehensive view of each component of the climate system and its changes to date. New climate model simulations, new analyses, and methods combining multiple lines of evidence lead to improved understanding of human influence on a wider range of climate variables, including weather and climate extremes. The time periods considered throughout this Section depend upon the availability of observational products, paleoclimate archives and peer-reviewed studies.

A.1 It is unequivocal that human influence has warmed the atmosphere, ocean and land. Widespread and rapid changes in the atmosphere, ocean, cryosphere and biosphere have occurred.

{2.2, 2.3, Cross-Chapter Box 2.3, 3.3, 3.4, 3.5, 3.6, 3.8, 5.2, 5.3, 6.4, 7.3, 8.3, 9.2, 9.3, 9.5, 9.6, Cross-Chapter Box 9.1} (Figure SPM.1, Figure SPM.2)

A.1.1 Observed increases in well-mixed greenhouse gas (GHG) concentrations since around 1750 are unequivocally caused by human activities. Since 2011 (measurements reported in AR5), concentrations have continued to increase in the atmosphere, reaching annual averages of 410 ppm for carbon dioxide (CO₂), 1866 ppb for methane (CH₄), and 332 ppb for nitrous oxide (N₂O) in 2019⁶. Land and ocean have taken up a near-constant proportion (globally about 56% per year) of CO₂ emissions from human activities over the past six decades, with regional differences (*high confidence*)⁷. {2.2, 5.2, 7.3, TS.2.2, Box TS.5}

A.1.2 Each of the last four decades has been successively warmer than any decade that preceded it since 1850. Global surface temperature⁸ in the first two decades of the 21st century (2001–2020) was 0.99 [0.84–1.10] °C higher than 1850–1900⁹. Global surface temperature was 1.09 [0.95 to 1.20] °C higher in 2011–2020 than 1850–1900, with larger increases over land (1.59 [1.34 to 1.83] °C) than over the ocean (0.88 [0.68 to 1.01] °C). The estimated increase in global surface temperature since AR5 is principally due to further warming since 2003–2012 (+0.19 [0.16 to 0.22] °C). Additionally, methodological advances and new datasets contributed approximately 0.1°C to the updated estimate of warming in AR6¹⁰.

⁶ Other GHG concentrations in 2019 were: PFCs (109 ppt CF₄ equivalent); SF₆ (10 ppt); NF₃ (2 ppt); HFCs (237 ppt HFC-134a equivalent); other Montreal Protocol gases (mainly CFCs, HCFCs, 1032 ppt CFC-12 equivalent). Increases from 2011 are 19 ppm for CO₂, 63 ppb for CH₄ and 8 ppb for N₂O.

⁷ Land and ocean are not substantial sinks for other GHGs.

⁸ The term ‘global surface temperature’ is used in reference to both global mean surface temperature and global surface air temperature throughout this SPM. Changes in these quantities are assessed with *high confidence* to differ by at most 10% from one another, but conflicting lines of evidence lead to *low confidence* in the sign of any difference in long-term trend. {Cross-Section Box TS.1}

⁹ The period 1850–1900 represents the earliest period of sufficiently globally complete observations to estimate global surface temperature and, consistent with AR5 and SR1.5, is used as an approximation for pre-industrial conditions.

¹⁰ Since AR5, methodological advances and new datasets have provided a more complete spatial representation of changes in surface temperature, including in the Arctic. These and other improvements have additionally increased the estimate of global surface temperature change by approximately 0.1 °C, but this increase does not represent additional physical warming since the AR5.

A.1.3 The *likely* range of total human-caused global surface temperature increase from 1850–1900 to 2010–2019¹¹ is 0.8°C to 1.3°C, with a best estimate of 1.07°C. It is *likely* that well-mixed GHGs contributed a warming of 1.0°C to 2.0°C, other human drivers (principally aerosols) contributed a cooling of 0.0°C to 0.8°C, natural drivers changed global surface temperature by –0.1°C to 0.1°C, and internal variability changed it by –0.2°C to 0.2°C. It is *very likely* that well-mixed GHGs were the main driver¹² of tropospheric warming since 1979, and *extremely likely* that human-caused stratospheric ozone depletion was the main driver of cooling of the lower stratosphere between 1979 and the mid-1990s. {3.3, 6.4, 7.3, Cross-Section Box TS.1, TS.2.3} (Figure SPM.2)

A.1.4 Globally averaged precipitation over land has *likely* increased since 1950, with a faster rate of increase since the 1980s (*medium confidence*). It is *likely* that human influence contributed to the pattern of observed precipitation changes since the mid-20th century, and *extremely likely* that human influence contributed to the pattern of observed changes in near-surface ocean salinity. Mid-latitude storm tracks have *likely* shifted poleward in both hemispheres since the 1980s, with marked seasonality in trends (*medium confidence*). For the Southern Hemisphere, human influence *very likely* contributed to the poleward shift of the closely related extratropical jet in austral summer. {2.3, 3.3, 8.3, 9.2, TS.2.3, TS.2.4, Box TS.6}

A.1.5 Human influence is *very likely* the main driver of the global retreat of glaciers since the 1990s and the decrease in Arctic sea ice area between 1979–1988 and 2010–2019 (about 40% in September and about 10% in March). There has been no significant trend in Antarctic sea ice area from 1979 to 2020 due to regionally opposing trends and large internal variability. Human influence *very likely* contributed to the decrease in Northern Hemisphere spring snow cover since 1950. It is *very likely* that human influence has contributed to the observed surface melting of the Greenland Ice Sheet over the past two decades, but there is only *limited evidence*, with *medium agreement*, of human influence on the Antarctic Ice Sheet mass loss. {2.3, 3.4, 8.3, 9.3, 9.5, TS.2.5}

A.1.6 It is *virtually certain* that the global upper ocean (0–700 m) has warmed since the 1970s and *extremely likely* that human influence is the main driver. It is *virtually certain* that human-caused CO₂ emissions are the main driver of current global acidification of the surface open ocean. There is *high confidence* that oxygen levels have dropped in many upper ocean regions since the mid-20th century, and *medium confidence* that human influence contributed to this drop. {2.3, 3.5, 3.6, 5.3, 9.2, TS.2.4}

A.1.7 Global mean sea level increased by 0.20 [0.15 to 0.25] m between 1901 and 2018. The average rate of sea level rise was 1.3 [0.6 to 2.1] mm yr⁻¹ between 1901 and 1971, increasing to 1.9 [0.8 to 2.9] mm yr⁻¹ between 1971 and 2006, and further increasing to 3.7 [3.2 to 4.2] mm yr⁻¹ between 2006 and 2018 (*high confidence*). Human influence was *very likely* the main driver of these increases since at least 1971. {2.3, 3.5, 9.6, Cross-Chapter Box 9.1, Box TS.4}

A.1.8 Changes in the land biosphere since 1970 are consistent with global warming: climate zones have shifted poleward in both hemispheres, and the growing season has on average lengthened by up to two days per decade since the 1950s in the Northern Hemisphere extratropics (*high confidence*). {2.3, TS.2.6}

¹¹ The period distinction with A.1.2 arises because the attribution studies consider this slightly earlier period. The observed warming to 2010–2019 is 1.06 [0.88 to 1.21] °C.


¹² Throughout this SPM, ‘main driver’ means responsible for more than 50% of the change.

Human influence has warmed the climate at a rate that is unprecedented in at least the last 2000 years

Attachment 1

Changes in global surface temperature relative to 1850-1900

a) Change in global surface temperature (decadal average) as reconstructed (1-2000) and observed (1850-2020)



b) Change in global surface temperature (annual average) as observed and simulated using human & natural and only natural factors (both 1850-2020)




Figure SPM.1: History of global temperature change and causes of recent warming.

Panel a): Changes in global surface temperature reconstructed from paleoclimate archives (solid grey line, 1–2000) and from direct observations (solid black line, 1850–2020), both relative to 1850–1900 and decadal averaged. The vertical bar on the left shows the estimated temperature (*very likely* range) during the warmest multi-century period in at least the last 100,000 years, which occurred around 6500 years ago during the current interglacial period (Holocene). The Last Interglacial, around 125,000 years ago, is the next most recent candidate for a period of higher temperature. These past warm periods were caused by slow (multi-millennial) orbital variations. The grey shading with white diagonal lines shows the *very likely* ranges for the temperature reconstructions.

Panel b): Changes in global surface temperature over the past 170 years (black line) relative to 1850–1900 and annually averaged, compared to CMIP6 climate model simulations (see Box SPM.1) of the temperature response to both human and natural drivers (brown), and to only natural drivers (solar and volcanic activity, green). Solid coloured lines show the multi-model average, and coloured shades show the *very likely* range of simulations. (see Figure SPM.2 for the assessed contributions to warming).

{2.3.1, 3.3, Cross-Chapter Box 2.3, Cross-Section Box TS.1, Figure 1a, TS.2.2}

Observed warming is driven by emissions from human activities, with greenhouse gas warming partly masked by aerosol cooling

Attachment 1




Figure SPM.2: Assessed contributions to observed warming in 2010–2019 relative to 1850–1900.

Panel a): Observed global warming (increase in global surface temperature) and its *very likely* range {3.3.1, Cross-Chapter Box 2.3}.

Panel b): Evidence from attribution studies, which synthesize information from climate models and observations. The panel shows temperature change attributed to total human influence, changes in well-mixed greenhouse gas concentrations, other human drivers due to aerosols, ozone and land-use change (land-use reflectance), solar and volcanic drivers, and internal climate variability. Whiskers show *likely* ranges {3.3.1}.

Panel c): Evidence from the assessment of radiative forcing and climate sensitivity. The panel shows temperature changes from individual components of human influence, including emissions of greenhouse gases, aerosols and their precursors; land-use changes (land-use reflectance and irrigation); and aviation contrails. Whiskers show *very likely* ranges. Estimates account for both direct emissions into the atmosphere and their effect, if any, on other climate drivers. For aerosols, both direct (through radiation) and indirect (through interactions with clouds) effects are considered. {6.4.2, 7.3}

A.2 The scale of recent changes across the climate system as a whole and the present state of many aspects of the climate system are unprecedented over many centuries to many thousands of years.

Attachment 1

{Cross-Chapter Box 2.1, 2.2, 2.3, 5.1} (Figure SPM.1)

A.2.1 In 2019, atmospheric CO₂ concentrations were higher than at any time in at least 2 million years (*high confidence*), and concentrations of CH₄ and N₂O were higher than at any time in at least 800,000 years (*very high confidence*). Since 1750, increases in CO₂ (47%) and CH₄ (156%) concentrations far exceed, and increases in N₂O (23%) are similar to, the natural multi-millennial changes between glacial and interglacial periods over at least the past 800,000 years (*very high confidence*).
{2.2, 5.1, TS.2.2}

A.2.2 Global surface temperature has increased faster since 1970 than in any other 50-year period over at least the last 2000 years (*high confidence*). Temperatures during the most recent decade (2011–2020) exceed those of the most recent multi-century warm period, around 6500 years ago¹³ [0.2°C to 1°C relative to 1850–1900] (*medium confidence*). Prior to that, the next most recent warm period was about 125,000 years ago when the multi-century temperature [0.5°C to 1.5°C relative to 1850–1900] overlaps the observations of the most recent decade (*medium confidence*).
{Cross-Chapter Box 2.1, 2.3, Cross-Section Box TS.1} (Figure SPM.1)

A.2.3 In 2011–2020, annual average Arctic sea ice area reached its lowest level since at least 1850 (*high confidence*). Late summer Arctic sea ice area was smaller than at any time in at least the past 1000 years (*medium confidence*). The global nature of glacier retreat, with almost all of the world's glaciers retreating synchronously, since the 1950s is unprecedented in at least the last 2000 years (*medium confidence*).
{2.3, TS.2.5}

A.2.4 Global mean sea level has risen faster since 1900 than over any preceding century in at least the last 3000 years (*high confidence*). The global ocean has warmed faster over the past century than since the end of the last deglacial transition (around 11,000 years ago) (*medium confidence*). A long-term increase in surface open ocean pH occurred over the past 50 million years (*high confidence*), and surface open ocean pH as low as recent decades is unusual in the last 2 million years (*medium confidence*).
{2.3, TS.2.4, Box TS.4}

¹³ As stated in section B.1, even under the very low emissions scenario SSP1-1.9, temperatures are assessed to remain elevated above those of the most recent decade until at least 2100 and therefore warmer than the century-scale period 6500 years ago.

A.3 Human-induced climate change is already affecting many weather and climate extremes in every region across the globe. Evidence of observed changes in extremes such as heatwaves, heavy precipitation, droughts, and tropical cyclones, and, in particular, their attribution to human influence, has strengthened since AR5. Attachment 1
 {2.3, 3.3, 8.2, 8.3, 8.4, 8.5, 8.6, Box 8.1, Box 8.2, Box 9.2, 10.6, 11.2, 11.3, 11.4, 11.6, 11.7, 11.8, 11.9, 12.3} (Figure SPM.3)

A.3.1 It is *virtually certain* that hot extremes (including heatwaves) have become more frequent and more intense across most land regions since the 1950s, while cold extremes (including cold waves) have become less frequent and less severe, with *high confidence* that human-induced climate change is the main driver¹⁴ of these changes. Some recent hot extremes observed over the past decade would have been *extremely unlikely* to occur without human influence on the climate system. Marine heatwaves have approximately doubled in frequency since the 1980s (*high confidence*), and human influence has *very likely* contributed to most of them since at least 2006.
 {Box 9.2, 11.2, 11.3, 11.9, TS.2.4, TS.2.6, Box TS.10} (Figure SPM.3)

A.3.2 The frequency and intensity of heavy precipitation events have increased since the 1950s over most land area for which observational data are sufficient for trend analysis (*high confidence*), and human-induced climate change is *likely* the main driver. Human-induced climate change has contributed to increases in agricultural and ecological droughts¹⁵ in some regions due to increased land evapotranspiration¹⁶ (*medium confidence*).
 {8.2, 8.3, 11.4, 11.6, 11.9, TS.2.6, Box TS.10} (Figure SPM.3)

A.3.3 Decreases in global land monsoon precipitation¹⁷ from the 1950s to the 1980s are partly attributed to human-caused Northern Hemisphere aerosol emissions, but increases since then have resulted from rising GHG concentrations and decadal to multi-decadal internal variability (*medium confidence*). Over South Asia, East Asia and West Africa increases in monsoon precipitation due to warming from GHG emissions were counteracted by decreases in monsoon precipitation due to cooling from human-caused aerosol emissions over the 20th century (*high confidence*). Increases in West African monsoon precipitation since the 1980s are partly due to the growing influence of GHGs and reductions in the cooling effect of human-caused aerosol emissions over Europe and North America (*medium confidence*).
 {2.3, 3.3, 8.2, 8.3, 8.4, 8.5, 8.6, Box 8.1, Box 8.2, 10.6, Box TS.13}

¹⁴ Throughout this SPM, ‘main driver’ means responsible for more than 50% of the change.

¹⁵ Agricultural and ecological drought (depending on the affected biome): a period with abnormal soil moisture deficit, which results from combined shortage of precipitation and excess evapotranspiration, and during the growing season impinges on crop production or ecosystem function in general. Observed changes in meteorological droughts (precipitation deficits) and hydrological droughts (streamflow deficits) are distinct from those in agricultural and ecological droughts and addressed in the underlying AR6 material (Chapter 11).

¹⁶ The combined processes through which water is transferred to the atmosphere from open water and ice surfaces, bare soil, and vegetation that make up the Earth’s surface.

¹⁷ The global monsoon is defined as the area in which the annual range (local summer minus local winter) of precipitation is greater than 2.5 mm day⁻¹. Global land monsoon precipitation refers to the mean precipitation over land areas within the global monsoon.

A.3.4 It is *likely* that the global proportion of major (Category 3–5) tropical cyclone occurrence has increased over the last four decades, and the latitude where tropical cyclones in the western North Pacific reach their peak intensity has shifted northward; these changes cannot be explained by internal variability alone (*medium confidence*). There is *low confidence* in long-term (multi-decadal to centennial) trends in the frequency of all-category tropical cyclones. Event attribution studies and physical understanding indicate that human-induced climate change increases heavy precipitation associated with tropical cyclones (*high confidence*) but data limitations inhibit clear detection of past trends on the global scale. {8.2, 11.7, Box TS.10} Attachment 1



A.3.5 Human influence has *likely* increased the chance of compound extreme events¹⁸ since the 1950s. This includes increases in the frequency of concurrent heatwaves and droughts on the global scale (*high confidence*); fire weather in some regions of all inhabited continents (*medium confidence*); and compound flooding in some locations (*medium confidence*). {11.6, 11.7, 11.8, 12.3, 12.4, TS.2.6, Table TS.5, Box TS.10}

¹⁸ Compound extreme events are the combination of multiple drivers and/or hazards that contribute to societal or environmental risk. Examples are concurrent heatwaves and droughts, compound flooding (e.g., a storm surge in combination with extreme rainfall and/or river flow), compound fire weather conditions (i.e., a combination of hot, dry, and windy conditions), or concurrent extremes at different locations.



Climate change is already affecting every inhabited region across the globe with human influence contributing to many observed changes in weather and climate extremes

Attachment 1



a) Synthesis of assessment of observed change in **hot extremes** and confidence in human contribution to the observed changes in the world's regions



b) Synthesis of assessment of observed change in **heavy precipitation** and confidence in human contribution to the observed changes in the world's regions



c) Synthesis of assessment of observed change in **agricultural and ecological drought** and confidence in human contribution to the observed changes in the world's regions



Each hexagon corresponds to one of the IPCC AR6 WGI reference regions

North-Western North America

IPCC AR6 WGI reference regions: **North America:** NWN (North-Western North America), NEN (North-Eastern North America), WNA (Western North America), CNA (Central North America), ENA (Eastern North America), **Central America:** NCA (Northern Central America), SCA (Southern Central America), CAR (Caribbean), **South America:** NWS (North-Western South America), NSA (Northern South America), NES (North-Eastern South America), SAM (South American Monsoon), SWS (South-Western South America), SES (South-Eastern South America), SSA (Southern South America), **Europe:** GIC (Greenland/Iceland), NEU (Northern Europe), WCE (Western and Central Europe), EEU (Eastern Europe), MED (Mediterranean), **Africa:** MED (Mediterranean), SAH (Sahara), WAF (Western Africa), CAF (Central Africa), NEAF (North Eastern Africa), SEAF (South Eastern Africa), WSAF (West Southern Africa), ESAF (East Southern Africa), MDG (Madagascar), **Asia:** RAR (Russian Arctic), WSB (West Siberia), ESB (East Siberia), RFE (Russian Far East), WCA (West Central Asia), ECA (East Central Asia), TIB (Tibetan Plateau), EAS (East Asia), ARP (Arabian Peninsula), SAS (South Asia), SEA (South East Asia), **Australasia:** NAU (Northern Australia), CAU (Central Australia), EAU (Eastern Australia), SAU (Southern Australia), NZ (New Zealand), **Small Islands:** CAR (Caribbean), PAC (Pacific Small Islands)

Figure SPM.3: Synthesis of assessed observed and attributable regional changes.

The IPCC AR6 WGI inhabited regions are displayed as **hexagons** with identical size in their approximate geographical location (see legend for regional acronyms). All assessments are made for each region as a whole and for the 1950s to the present. Assessments made on different time scales or more local spatial scales might differ from what is shown in the figure. The **colours** in each panel represent the four outcomes of the assessment on observed changes. White and light grey striped hexagons are used where there is *low agreement* in the type of change for the region as a whole, and grey hexagons are used when there is limited data and/or literature that prevents an assessment of the region as a whole. Other colours indicate at least *medium confidence* in the observed change. The **confidence level** for the human influence on these observed changes is based on assessing trend detection and attribution and event attribution literature, and it is indicated by the number of dots: three dots for *high confidence*, two dots for *medium confidence* and one dot for *low confidence* (filled: limited agreement; empty: limited evidence).

Panel a) For hot extremes, the evidence is mostly drawn from changes in metrics based on daily maximum temperatures; regional studies using other indices (heatwave duration, frequency and intensity) are used in addition. Red hexagons indicate regions where there is at least *medium confidence* in an observed increase in hot extremes.

Panel b) For heavy precipitation, the evidence is mostly drawn from changes in indices based on one-day or five-day precipitation amounts using global and regional studies. Green hexagons indicate regions where there is at least *medium confidence* in an observed increase in heavy precipitation.

Panel c) Agricultural and ecological droughts are assessed based on observed and simulated changes in total column soil moisture, complemented by evidence on changes in surface soil moisture, water balance (precipitation minus evapotranspiration) and indices driven by precipitation and atmospheric evaporative demand. Yellow hexagons indicate regions where there is at least *medium confidence* in an observed increase in this type of drought and green hexagons indicate regions where there is at least *medium confidence* in an observed decrease in agricultural and ecological drought.

For all regions, table TS.5 shows a broader range of observed changes besides the ones shown in this figure. Note that SSA is the only region that does not display observed changes in the metrics shown in this figure, but is affected by observed increases in mean temperature, decreases in frost, and increases in marine heatwaves.

{11.9, Table TS.5, Box TS.10, Figure 1, Atlas 1.3.3, Figure Atlas.2}

A.4 Improved knowledge of climate processes, paleoclimate evidence and the response of the climate system to increasing radiative forcing gives a best estimate of equilibrium climate sensitivity of 3°C with a narrower range compared to AR5. {2.2, 7.3, 7.4, 7.5, Box 7.2, Cross-Chapter Box 9.1, 9.4, 9.5, 9.6}

A.4.1 Human-caused radiative forcing of 2.72 [1.96 to 3.48] W m⁻² in 2019 relative to 1750 has warmed the climate system. This warming is mainly due to increased GHG concentrations, partly reduced by cooling due to increased aerosol concentrations. The radiative forcing has increased by 0.43 W m⁻² (19%) relative to AR5, of which 0.34 W m⁻² is due to the increase in GHG concentrations since 2011. The remainder is due to improved scientific understanding and changes in the assessment of aerosol forcing, which include decreases in concentration and improvement in its calculation (*high confidence*).

{2.2, 7.3, TS.2.2, TS.3.1}

A.4.2 Human-caused net positive radiative forcing causes an accumulation of additional energy (heating) in the climate system, partly reduced by increased energy loss to space in response to surface warming. The observed average rate of heating of the climate system increased from 0.50 [0.32 to 0.69] W m^{-2} for the period 1971–2006¹⁹, to 0.79 [0.52 to 1.06] W m^{-2} for the period 2006–2018²⁰ (*high confidence*). Ocean warming accounted for 91% of the heating in the climate system, with land warming, ice loss and atmospheric warming accounting for about 5%, 3% and 1%, respectively (*high confidence*).
{7.2, Box 7.2, TS.3.1}

A.4.3 Heating of the climate system has caused global mean sea level rise through ice loss on land and thermal expansion from ocean warming. Thermal expansion explained 50% of sea level rise during 1971–2018, while ice loss from glaciers contributed 22%, ice sheets 20% and changes in land water storage 8%. The rate of ice sheet loss increased by a factor of four between 1992–1999 and 2010–2019. Together, ice sheet and glacier mass loss were the dominant contributors to global mean sea level rise during 2006–2018. (*high confidence*)
{Cross-Chapter Box 9.1, 9.4, 9.5, 9.6}

A.4.4 The equilibrium climate sensitivity is an important quantity used to estimate how the climate responds to radiative forcing. Based on multiple lines of evidence²¹, the *very likely* range of equilibrium climate sensitivity is between 2°C (*high confidence*) and 5°C (*medium confidence*). The AR6 assessed best estimate is 3°C with a *likely* range of 2.5°C to 4°C (*high confidence*), compared to 1.5°C to 4.5°C in AR5, which did not provide a best estimate.
{7.4, 7.5, TS.3.2}

¹⁹ cumulative energy increase of 282 [177 to 387] ZJ over 1971–2006 (1 ZJ = 10^{21} J).

²⁰ cumulative energy increase of 152 [100 to 205] ZJ over 2006–2018.

²¹ Understanding of climate processes, the instrumental record, paleoclimates and model-based emergent constraints (see glossary).

B. Possible Climate Futures

Attachment 1

A set of five new illustrative emissions scenarios is considered consistently across this report to explore the climate response to a broader range of greenhouse gas (GHG), land use and air pollutant futures than assessed in AR5. This set of scenarios drives climate model projections of changes in the climate system. These projections account for solar activity and background forcing from volcanoes. Results over the 21st century are provided for the near-term (2021–2040), mid-term (2041–2060) and long-term (2081–2100) relative to 1850–1900, unless otherwise stated.

Box SPM.1: Scenarios, Climate Models and Projections

Box SPM.1.1: This report assesses the climate response to five illustrative scenarios that cover the range of possible future development of anthropogenic drivers of climate change found in the literature. They start in 2015, and include scenarios²² with high and very high GHG emissions (SSP3-7.0 and SSP5-8.5) and CO₂ emissions that roughly double from current levels by 2100 and 2050, respectively, scenarios with intermediate GHG emissions (SSP2-4.5) and CO₂ emissions remaining around current levels until the middle of the century, and scenarios with very low and low GHG emissions and CO₂ emissions declining to net zero around or after 2050, followed by varying levels of net negative CO₂ emissions²³ (SSP1-1.9 and SSP1-2.6) as illustrated in Figure SPM.4. Emissions vary between scenarios depending on socio-economic assumptions, levels of climate change mitigation and, for aerosols and non-methane ozone precursors, air pollution controls. Alternative assumptions may result in similar emissions and climate responses, but the socio-economic assumptions and the feasibility or likelihood of individual scenarios is not part of the assessment.

{TS.1.3, 1.6, Cross-Chapter Box 1.4} (Figure SPM.4)

Box SPM.1.2: This report assesses results from climate models participating in the Coupled Model Intercomparison Project Phase 6 (CMIP6) of the World Climate Research Programme. These models include new and better representation of physical, chemical and biological processes, as well as higher resolution, compared to climate models considered in previous IPCC assessment reports. This has improved the simulation of the recent mean state of most large-scale indicators of climate change and many other aspects across the climate system. Some differences from observations remain, for example in regional precipitation patterns. The CMIP6 historical simulations assessed in this report have an ensemble mean global surface temperature change within 0.2°C of the observations over most of the historical period, and observed warming is within the *very likely* range of the CMIP6 ensemble. However, some CMIP6 models simulate a warming that is either above or below the assessed *very likely* range of observed warming.

{1.5, Cross-Chapter Box 2.2, 3.3, 3.8, TS.1.2, Cross-Section Box TS.1} (Figure SPM.1 b, Figure SPM.2)

Box SPM.1.3: The CMIP6 models considered in this Report have a wider range of climate sensitivity than in CMIP5 models and the AR6 assessed *very likely* range, which is based on multiple lines of evidence. These CMIP6 models also show a higher average climate sensitivity than CMIP5 and the AR6 assessed best estimate. The higher CMIP6 climate sensitivity values compared to CMIP5 can be traced to an amplifying cloud feedback that is larger in CMIP6 by about 20%.

{Box 7.1, 7.3, 7.4, 7.5, TS.3.2}

Box SPM.1.4: For the first time in an IPCC report, assessed future changes in global surface temperature, ocean warming and sea level are constructed by combining multi-model projections with observational constraints based on past simulated warming, as well as the AR6 assessment of climate sensitivity. For other quantities, such robust methods do not yet exist to constrain the projections. Nevertheless, robust projected

²² Throughout this report, the five illustrative scenarios are referred to as SSPx-y, where ‘SSPx’ refers to the Shared Socio-economic Pathway or ‘SSP’ describing the socio-economic trends underlying the scenario, and ‘y’ refers to the approximate level of radiative forcing (in W m⁻²) resulting from the scenario in the year 2100. A detailed comparison to scenarios used in earlier IPCC reports is provided in Section TS.1.3 and 1.6 and 4.6. The SSPs that underlie the specific forcing scenarios used to drive climate models are not assessed by WGI. Rather, the SSPx-y labelling ensures traceability to the underlying literature in which specific forcing pathways are used as input to the climate models. IPCC is neutral with regard to the assumptions underlying the SSPs, which do not cover all possible scenarios. Alternative scenarios may be considered or developed.


²³ Net negative CO₂ emissions are reached when anthropogenic removals of CO₂ exceed anthropogenic emissions. {Glossary}

geographical patterns of many variables can be identified at a given level of global warming, common to all scenarios considered and independent of timing when the global warming level is reached. {1.6, Box 4.1, 4.3, 4.6, 7.5, 9.2, 9.6, Cross-Chapter Box 11.1, Cross-Section Box TS.1}

Attachment 1


Future emissions cause future additional warming, with total warming dominated by past and future CO₂ emissions

a) Future annual emissions of CO₂ (left) and of a subset of key non-CO₂ drivers (right), across five illustrative scenarios



b) Contribution to global surface temperature increase from different emissions, with a dominant role of CO₂ emissions

Change in global surface temperature in 2081-2100 relative to 1850-1900 (°C)



Total warming (observed warming to date in darker shade), warming from CO₂, warming from non-CO₂ GHGs and cooling from changes in aerosols and land use

Figure SPM.4: Future anthropogenic emissions of key drivers of climate change and warming contributions by groups of drivers for the five illustrative scenarios used in this report.

Attachment 1

The five scenarios are SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP3-7.0 and SSP5-8.5.

Panel a) Annual anthropogenic (human-caused) emissions over the 2015–2100 period. Shown are emissions trajectories for carbon dioxide (CO₂) from all sectors (GtCO₂/yr) (left graph) and for a subset of three key non-CO₂ drivers considered in the scenarios: methane (CH₄, MtCH₄/yr, top-right graph), nitrous oxide (N₂O, MtN₂O/yr, middle-right graph) and sulfur dioxide (SO₂, MtSO₂/yr, bottom-right graph, contributing to anthropogenic aerosols in panel b).

Panel b) Warming contributions by groups of anthropogenic drivers and by scenario are shown as change in global surface temperature (°C) in 2081–2100 relative to 1850–1900, with indication of the observed warming to date. Bars and whiskers represent median values and the *very likely* range, respectively. Within each scenario bar plot, the bars represent total global warming (°C; total bar) (see Table SPM.1) and warming contributions (°C) from changes in CO₂ (CO₂ bar), from non-CO₂ greenhouse gases (non-CO₂ GHGs bar; comprising well-mixed greenhouse gases and ozone) and net cooling from other anthropogenic drivers (aerosols and land-use bar; anthropogenic aerosols, changes in reflectance due to land-use and irrigation changes, and contrails from aviation; see Figure SPM.2, panel c, for the warming contributions to date for individual drivers). The best estimate for observed warming in 2010–2019 relative to 1850–1900 (see Figure SPM.2, panel a) is indicated in the darker column in the total bar. Warming contributions in panel b are calculated as explained in Table SPM.1 for the total bar. For the other bars the contribution by groups of drivers are calculated with a physical climate emulator of global surface temperature which relies on climate sensitivity and radiative forcing assessments.

{Cross-Chapter Box 1.4, 4.6, Figure 4.35, 6.7, Figure 6.18, 6.22 and 6.24, Cross-Chapter Box 7.1, 7.3, Figure 7.7, Box TS.7, Figures TS.4 and TS.15}

B.1 Global surface temperature will continue to increase until at least the mid-century under all emissions scenarios considered. Global warming of 1.5°C and 2°C will be exceeded during the 21st century unless deep reductions in CO₂ and other greenhouse gas emissions occur in the coming decades.
{2.3, Cross-Chapter Box 2.3, Cross-Chapter Box 2.4, 4.3, 4.4, 4.5} (Figure SPM.1, Figure SPM.4, Figure SPM.8, Table SPM.1, Box SPM.1)

B.1.1 Compared to 1850–1900, global surface temperature averaged over 2081–2100 is *very likely* to be higher by 1.0°C to 1.8°C under the very low GHG emissions scenario considered (SSP1-1.9), by 2.1°C to 3.5°C in the intermediate scenario (SSP2-4.5) and by 3.3°C to 5.7°C under the very high GHG emissions scenario (SSP5-8.5)²⁴. The last time global surface temperature was sustained at or above 2.5°C higher than 1850–1900 was over 3 million years ago (*medium confidence*).
{2.3, Cross-Chapter Box 2.4, 4.3, 4.5, Box TS.2, Box TS.4, Cross-Section Box TS.1} (Table SPM.1)

Table SPM.1: Changes in global surface temperature, which are assessed based on multiple lines of evidence, for selected 20-year time periods and the five illustrative emissions scenarios considered. Temperature differences relative to the average global surface temperature of the period 1850–1900 are reported in °C. This includes the revised assessment of observed historical warming for the AR5 reference period 1986–2005, which in AR6 is higher by 0.08 [–0.01 to 0.12] °C than in the AR5 (see footnote 10). Changes relative to the recent reference period 1995–2014 may be calculated approximately by subtracting 0.85°C, the best estimate of the observed warming from 1850–1900 to 1995–2014.
{Cross-Chapter Box 2.3, 4.3, 4.4, Cross-Section Box TS.1}

²⁴ Changes in global surface temperature are reported as running 20-year averages, unless stated otherwise.

Scenario	Near term, 2021–2040		Mid-term, 2041–2060		Long term, 2081–2100	
	Best estimate (°C)	Very likely range (°C)	Best estimate (°C)	Very likely range (°C)	Best estimate (°C)	Very likely range (°C)
SSP1-1.9	1.5	1.2 to 1.7	1.6	1.2 to 2.0	1.4	1.0 to 1.8
SSP1-2.6	1.5	1.2 to 1.8	1.7	1.3 to 2.2	1.8	1.3 to 2.4
SSP2-4.5	1.5	1.2 to 1.8	2.0	1.6 to 2.5	2.7	2.1 to 3.5
SSP3-7.0	1.5	1.2 to 1.8	2.1	1.7 to 2.6	3.6	2.8 to 4.6
SSP5-8.5	1.6	1.3 to 1.9	2.4	1.9 to 3.0	4.4	3.3 to 5.7

B.1.2 Based on the assessment of multiple lines of evidence, global warming of 2°C, relative to 1850–1900, would be exceeded during the 21st century under the high and very high GHG emissions scenarios considered in this report (SSP3-7.0 and SSP5-8.5, respectively). Global warming of 2°C would *extremely likely* be exceeded in the intermediate scenario (SSP2-4.5). Under the very low and low GHG emissions scenarios, global warming of 2°C is *extremely unlikely* to be exceeded (SSP1-1.9), or *unlikely* to be exceeded (SSP1-2.6)²⁵. Crossing the 2°C global warming level in the mid-term period (2041–2060) is *very likely* to occur under the very high GHG emissions scenario (SSP5-8.5), *likely* to occur under the high GHG emissions scenario (SSP3-7.0), and *more likely than not* to occur in the intermediate GHG emissions scenario (SSP2-4.5)²⁶.

{4.3, Cross-Section Box TS.1} (Table SPM.1, Figure SPM.4, Box SPM.1)

B.1.3 Global warming of 1.5°C relative to 1850-1900 would be exceeded during the 21st century under the intermediate, high and very high scenarios considered in this report (SSP2-4.5, SSP3-7.0 and SSP5-8.5, respectively). Under the five illustrative scenarios, in the near term (2021-2040), the 1.5°C global warming level is *very likely* to be exceeded under the very high GHG emissions scenario (SSP5-8.5), *likely* to be exceeded under the intermediate and high GHG emissions scenarios (SSP2-4.5 and SSP3-7.0), *more likely than not* to be exceeded under the low GHG emissions scenario (SSP1-2.6) and *more likely than not* to be reached under the very low GHG emissions scenario (SSP1-1.9)²⁷. Furthermore, for the very low GHG emissions scenario (SSP1-1.9), it is *more likely than not* that global surface temperature would decline back to below 1.5°C toward the end of the 21st century, with a temporary overshoot of no more than 0.1°C above 1.5°C global warming.

{4.3, Cross-Section Box TS.1} (Table SPM.1, Figure SPM.4)

²⁵ SSP1-1.9 and SSP1-2.6 are scenarios that start in 2015 and have very low and low GHG emissions and CO₂ emissions declining to net zero around or after 2050, followed by varying levels of net negative CO₂ emissions.

²⁶ Crossing is defined here as having the assessed global surface temperature change, averaged over a 20-year period, exceed a particular global warming level.

²⁷ The AR6 assessment of when a given global warming level is first exceeded benefits from the consideration of the illustrative scenarios, the multiple lines of evidence entering the assessment of future global surface temperature response to radiative forcing, and the improved estimate of historical warming. The AR6 assessment is thus not directly comparable to the SR1.5 SPM, which reported likely reaching 1.5°C global warming between 2030 and 2052, from a simple linear extrapolation of warming rates of the recent past. When considering scenarios similar to SSP1-1.9 instead of linear extrapolation, the SR1.5 estimate of when 1.5°C global

B.1.4 Global surface temperature in any single year can vary above or below the long-term human-induced trend, due to substantial natural variability²⁸. The occurrence of individual years with global surface temperature change above a certain level, for example 1.5°C or 2°C, relative to 1850–1900 does not imply that this global warming level has been reached²⁹. Attachment 1

{Cross-Chapter Box 2.3, 4.3, 4.4, Box 4.1, Cross-Section Box TS.1} (Table SPM.1, Figure SPM.1, Figure SPM.8)

B.2 Many changes in the climate system become larger in direct relation to increasing global warming. They include increases in the frequency and intensity of hot extremes, marine heatwaves, and heavy precipitation, agricultural and ecological droughts in some regions, and proportion of intense tropical cyclones, as well as reductions in Arctic sea ice, snow cover and permafrost. {4.3, 4.5, 4.6, 7.4, 8.2, 8.4, Box 8.2, 9.3, 9.5, Box 9.2, 11.1, 11.2, 11.3, 11.4, 11.6, 11.7, 11.9, Cross-Chapter Box 11.1, 12.4, 12.5, Cross-Chapter Box 12.1, Atlas.4, Atlas.5, Atlas.6, Atlas.7, Atlas.8, Atlas.9, Atlas.10, Atlas.11} (Figure SPM.5, Figure SPM.6, Figure SPM.8)

B.2.1 It is *virtually certain* that the land surface will continue to warm more than the ocean surface (*likely* 1.4 to 1.7 times more). It is *virtually certain* that the Arctic will continue to warm more than global surface temperature, with *high confidence* above two times the rate of global warming.

{2.3, 4.3, 4.5, 4.6, 7.4, 11.1, 11.3, 11.9, 12.4, 12.5, Cross-Chapter Box 12.1, Atlas.4, Atlas.5, Atlas.6, Atlas.7, Atlas.8, Atlas.9, Atlas.10, Atlas.11, Cross-Section Box TS.1, TS.2.6} (Figure SPM.5)

B.2.2 With every additional increment of global warming, changes in extremes continue to become larger. For example, every additional 0.5°C of global warming causes clearly discernible increases in the intensity and frequency of hot extremes, including heatwaves (*very likely*), and heavy precipitation (*high confidence*), as well as agricultural and ecological droughts³⁰ in some regions (*high confidence*). Discernible changes in intensity and frequency of meteorological droughts, with more regions showing increases than decreases, are seen in some regions for every additional 0.5°C of global warming (*medium confidence*). Increases in frequency and intensity of hydrological droughts become larger with increasing global warming in some regions (*medium confidence*). There will be an increasing occurrence of some extreme events unprecedented in the observational record with additional global warming, even at 1.5°C of global warming. Projected percentage changes in frequency are higher for rarer events (*high confidence*).

{8.2, 11.2, 11.3, 11.4, 11.6, 11.9, Cross-Chapter Box 11.1, Cross-Chapter Box 12.1, TS.2.6} (Figure SPM.5, Figure SPM.6)

warming is first exceeded is close to the best estimate reported here.

²⁸ Natural variability refers to climatic fluctuations that occur without any human influence, that is, internal variability combined with the response to external natural factors such as volcanic eruptions, changes in solar activity and, on longer time scales, orbital effects and plate tectonics.

²⁹ The internal variability in any single year is estimated to be $\pm 0.25^\circ\text{C}$ (5–95% range, *high confidence*).

³⁰ Projected changes in agricultural and ecological droughts are primarily assessed based on total column soil moisture. See footnote 15 for definition and relation to precipitation and evapotranspiration.

B.2.3 Some mid-latitude and semi-arid regions, and the South American Monsoon region, are projected to see the highest increase in the temperature of the hottest days, at about 1.5 to 2 times the rate of global warming (*high confidence*). The Arctic is projected to experience the highest increase in the temperature of the coldest days, at about 3 times the rate of global warming (*high confidence*). With additional global warming, the frequency of marine heatwaves will continue to increase (*high confidence*), particularly in the tropical ocean and the Arctic (*medium confidence*). Attachment 1

{Box 9.2, 11.1, 11.3, 11.9, Cross-Chapter Box 11.1, Cross-Chapter Box 12.1, 12.4, TS.2.4, TS.2.6} (**Figure SPM.6**)

B.2.4 It is *very likely* that heavy precipitation events will intensify and become more frequent in most regions with additional global warming. At the global scale, extreme daily precipitation events are projected to intensify by about 7% for each 1°C of global warming (*high confidence*). The proportion of intense tropical cyclones (categories 4-5) and peak wind speeds of the most intense tropical cyclones are projected to increase at the global scale with increasing global warming (*high confidence*).

{8.2, 11.4, 11.7, 11.9, Cross-Chapter Box 11.1, Box TS.6, TS.4.3.1} (**Figure SPM.5, Figure SPM.6**)

B.2.5 Additional warming is projected to further amplify permafrost thawing, and loss of seasonal snow cover, of land ice and of Arctic sea ice (*high confidence*). The Arctic is *likely* to be practically sea ice free in September³¹ at least once before 2050 under the five illustrative scenarios considered in this report, with more frequent occurrences for higher warming levels. There is *low confidence* in the projected decrease of Antarctic sea ice.

{4.3, 4.5, 7.4, 8.2, 8.4, Box 8.2, 9.3, 9.5, 12.4, Cross-Chapter Box 12.1, Atlas.5, Atlas.6, Atlas.8, Atlas.9, Atlas.11, TS.2.5} (**Figure SPM.8**)

³¹ monthly average sea ice area of less than 1 million km² which is about 15% of the average September sea ice area observed in 1979-1988

With every increment of global warming, changes get larger in regional mean temperature, precipitation and soil moisture

Attachment 1




Figure SPM.5: Changes in annual mean surface temperature, precipitation, and soil moisture.

Panel a) Comparison of observed and simulated annual mean surface temperature change. The left map shows the observed changes in annual mean surface temperature in the period of 1850–2020 per °C of global warming (°C). The local (i.e., grid point) observed annual mean surface temperature changes are linearly regressed against the global surface temperature in the period 1850–2020. Observed temperature data are from Berkeley Earth, the dataset with the largest coverage and highest horizontal resolution. Linear regression is applied to all years for which data at the corresponding grid point is available. The regression method was used to take into account the complete observational time series and thereby reduce the role of internal variability at the grid point level. White indicates areas where time coverage was 100 years or less and thereby too short to calculate a reliable linear regression. The **right map** is based on model simulations and shows change in annual multi-model mean simulated temperatures at a global warming level of 1°C (20-year mean global surface temperature change relative to 1850–1900). The triangles at each end of the color bar indicate out-of-bound values, that is, values above or below the given limits.

Panel b) Simulated annual mean temperature change (°C), panel c) precipitation change (%), and panel d) total column soil moisture change (standard deviation of interannual variability) at global warming levels of 1.5°C, 2°C and 4°C (20-yr mean global surface temperature change relative to 1850–1900). Simulated changes correspond to CMIP6 multi-model mean change (median change for soil moisture) at the corresponding global warming level, i.e. the same method as for the right map in panel a).




c) Annual mean precipitation change (%) relative to 1850-1900

Precipitation is projected to increase over high latitudes, the equatorial Pacific and parts of the monsoon regions, but decrease over parts of the subtropics and in limited areas of the tropics.


Simulated change at 1.5 °C global warming

Simulated change at 2 °C global warming

Simulated change at 4 °C global warming



Relatively small absolute changes may appear as large % changes in regions with dry baseline conditions






d) Annual mean total column soil moisture change (standard deviation)

Across warming levels, changes in soil moisture largely follow changes in precipitation but also show some differences due to the influence of evapotranspiration.


Simulated change at 1.5 °C global warming

Simulated change at 2 °C global warming

Simulated change at 4 °C global warming



Relatively small absolute changes may appear large when expressed in units of standard deviation in dry regions with little interannual variability in baseline conditions



In **panel c)**, high positive percentage changes in dry regions may correspond to small absolute changes. In **panel d)**, the unit is the standard deviation of interannual variability in soil moisture during 1850–1900. Standard deviation is a widely used metric in characterizing drought severity. A projected reduction in mean soil moisture by one standard deviation corresponds to soil moisture conditions typical of droughts that occurred about once every six years during 1850–1900. In panel d), large changes in dry regions with little interannual variability in the baseline conditions can correspond to small absolute change. The triangles at each end of the color bars indicate out-of-bound values, that is, values above or below the given limits. Results from all models reaching the corresponding warming level in any of the five illustrative scenarios (SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP3-7.0 and SSP5-8.5) are averaged. Maps of annual mean temperature and precipitation changes at a global warming level of 3°C are available in Figure 4.31 and Figure 4.32 in Section 4.6.

Corresponding maps of panels b), c) and d) including hatching to indicate the level of model agreement at grid-cell level are found in Figures 4.31, 4.32 and 11.19, respectively; as highlighted in CC-box Atlas.1, grid-cell level hatching is not informative for larger spatial scales (e.g., over AR6 reference regions) where the aggregated signals are less affected by small-scale variability leading to an increase in robustness.

{TS.1.3.2, Figure TS.3, Figure TS.5, Figure 1.14, 4.6.1, Cross-Chapter Box 11.1, Cross-Chapter Box Atlas.1}

Projected changes in extremes are larger in frequency and intensity with every additional increment of global warming

Attachment 1






Figure SPM.6: Projected changes in the intensity and frequency of hot temperature extremes over land, extreme precipitation over land, and agricultural and ecological droughts in drying regions Attachment 1

Projected changes are shown at global warming levels of 1°C, 1.5°C, 2°C, and 4°C and are relative to 1850–1900⁹ representing a climate without human influence. The figure depicts frequencies and increases in intensity of 10- or 50-year extreme events from the base period (1850–1900) under different global warming levels.

Hot temperature extremes are defined as the daily maximum temperatures over land that were exceeded on average once in a decade (10-year event) or once in 50 years (50-year event) during the 1850–1900 reference period. **Extreme precipitation events** are defined as the daily precipitation amount over land that was exceeded on average once in a decade during the 1850–1900 reference period. **Agricultural and ecological drought events** are defined as the annual average of total column soil moisture below the 10th percentile of the 1850–1900 base period. These extremes are defined on model grid box scale. For hot temperature extremes and extreme precipitation, results are shown for the global land. For agricultural and ecological drought, results are shown for drying regions only, which correspond to the AR6 regions in which there is at least *medium confidence* in a projected increase in agricultural/ecological drought at the 2°C warming level compared to the 1850–1900 base period in CMIP6. These regions include W. North-America, C. North-America, N. Central-America, S. Central-America, Caribbean, N. South-America, N.E. South-America, South-American-Monsoon, S.W. South-America, S. South-America, West & Central-Europe, Mediterranean, W. Southern-Africa, E. Southern-Africa, Madagascar, E. Australia, S. Australia (Caribbean is not included in the calculation of the figure because of the too small number of full land grid cells). The non-drying regions do not show an overall increase or decrease in drought severity. Projections of changes in agricultural and ecological droughts in the CMIP5 multi-model ensemble differ from those in CMIP6 in some regions, including in part of Africa and Asia. Assessments on projected changes in meteorological and hydrological droughts are provided in Chapter 11. {11.6, 11.9}

In the **‘frequency’ section**, each year is represented by a dot. The dark dots indicate years in which the extreme threshold is exceeded, while light dots are years when the threshold is not exceeded. Values correspond to the medians (in bold) and their respective 5–95% range based on the multi-model ensemble from simulations of CMIP6 under different SSP scenarios. For consistency, the number of dark dots is based on the rounded-up median. In the **‘intensity’ section**, medians and their 5–95% range, also based on the multi-model ensemble from simulations of CMIP6, are displayed as dark and light bars, respectively. Changes in the intensity of hot temperature extremes and extreme precipitations are expressed as degree Celsius and percentage. As for agricultural and ecological drought, intensity changes are expressed as fractions of standard deviation of annual soil moisture.

{11.1, 11.3, 11.4, 11.6, Figure 11.12, Figure 11.15, Figure 11.6, Figure 11.7, Figure 11.18}

B.3 Continued global warming is projected to further intensify the global water cycle, including its variability, global monsoon precipitation and the severity of wet and dry events.

Attachment 1

{4.3, 4.4, 4.5, 4.6, 8.2, 8.3, 8.4, 8.5, Box 8.2, 11.4, 11.6, 11.9, 12.4, Atlas.3} (Figure SPM.5, Figure SPM.6)

B.3.1 There is strengthened evidence since AR5 that the global water cycle will continue to intensify as global temperatures rise (*high confidence*), with precipitation and surface water flows projected to become more variable over most land regions within seasons (*high confidence*) and from year to year (*medium confidence*). The average annual global land precipitation is projected to increase by 0–5% under the very low GHG emissions scenario (SSP1-1.9), 1.5–8% for the intermediate GHG emissions scenario (SSP2-4.5) and 1–13% under the very high GHG emissions scenario (SSP5-8.5) by 2081–2100 relative to 1995–2014 (*likely* ranges). Precipitation is projected to increase over high latitudes, the equatorial Pacific and parts of the monsoon regions, but decrease over parts of the subtropics and limited areas in the tropics in SSP2-4.5, SSP3-7.0 and SSP5-8.5 (*very likely*). The portion of the global land experiencing detectable increases or decreases in seasonal mean precipitation is projected to increase (*medium confidence*). There is *high confidence* in an earlier onset of spring snowmelt, with higher peak flows at the expense of summer flows in snow-dominated regions globally.

{4.3, 4.5, 4.6, 8.2, 8.4, Atlas.3, TS.2.6, Box TS.6, TS.4.3} (Figure SPM.5)

B.3.2 A warmer climate will intensify very wet and very dry weather and climate events and seasons, with implications for flooding or drought (*high confidence*), but the location and frequency of these events depend on projected changes in regional atmospheric circulation, including monsoons and mid-latitude storm tracks. It is *very likely* that rainfall variability related to the El Niño–Southern Oscillation is projected to be amplified by the second half of the 21st century in the SSP2-4.5, SSP3-7.0 and SSP5-8.5 scenarios.

{4.3, 4.5, 4.6, 8.2, 8.4, 8.5, 11.4, 11.6, 11.9, 12.4, TS.2.6, TS.4.2, Box TS.6} (Figure SPM.5, Figure SPM.6)

B.3.3 Monsoon precipitation is projected to increase in the mid- to long term at global scale, particularly over South and Southeast Asia, East Asia and West Africa apart from the far west Sahel (*high confidence*). The monsoon season is projected to have a delayed onset over North and South America and West Africa (*high confidence*) and a delayed retreat over West Africa (*medium confidence*).

{4.4, 4.5, 8.2, 8.3, 8.4, Box 8.2, Box TS.13}

B.3.4 A projected southward shift and intensification of Southern Hemisphere summer mid-latitude storm tracks and associated precipitation is *likely* in the long term under high GHG emissions scenarios (SSP3-7.0, SSP5-8.5), but in the near term the effect of stratospheric ozone recovery counteracts these changes (*high confidence*). There is *medium confidence* in a continued poleward shift of storms and their precipitation in the North Pacific, while there is *low confidence* in projected changes in the North Atlantic storm tracks.

{TS.4.2, 4.4, 4.5, 8.4, TS.2.3}

B.4 Under scenarios with increasing CO₂ emissions, the ocean and land carbon sinks are projected to be less effective at slowing the accumulation of CO₂ in the atmosphere.

{4.3, 5.2, 5.4, 5.5, 5.6} (Figure SPM.7)

B.4.1 While natural land and ocean carbon sinks are projected to take up, in absolute terms, a progressively larger amount of CO₂ under higher compared to lower CO₂ emissions scenarios, they become less effective, that is, the proportion of emissions taken up by land and ocean decrease with increasing cumulative CO₂ emissions. This is projected to result in a higher proportion of emitted CO₂ remaining in the atmosphere (*high confidence*).

{5.2, 5.4, Box TS.5} (Figure SPM.7)

B.4.2 Based on model projections, under the intermediate scenario that stabilizes atmospheric CO₂ concentrations this century (SSP2-4.5), the rates of CO₂ taken up by the land and oceans are projected to decrease in the second half of the 21st century (*high confidence*). Under the very low and low GHG emissions scenarios (SSP1-1.9, SSP1-2.6), where CO₂ concentrations peak and decline during the 21st century, land and oceans begin to take up less carbon in response to declining atmospheric CO₂ concentrations (*high confidence*) and turn into a weak net source by 2100 under SSP1-1.9 (*medium confidence*). It is *very unlikely* that the combined global land and ocean sink will turn into a source by 2100 under scenarios without net negative emissions³² (SSP2-4.5, SSP3-7.0, SSP5-8.5).
{4.3, 5.4, 5.5, 5.6, Box TS.5, TS.3.3}

B.4.3 The magnitude of feedbacks between climate change and the carbon cycle becomes larger but also more uncertain in high CO₂ emissions scenarios (*very high confidence*). However, climate model projections show that the uncertainties in atmospheric CO₂ concentrations by 2100 are dominated by the differences between emissions scenarios (*high confidence*). Additional ecosystem responses to warming not yet fully included in climate models, such as CO₂ and CH₄ fluxes from wetlands, permafrost thaw and wildfires, would further increase concentrations of these gases in the atmosphere (*high confidence*).
{5.4, Box TS.5, TS.3.2}

³² These projected adjustments of carbon sinks to stabilization or decline of atmospheric CO₂ are accounted for in calculations of remaining carbon budgets.

The proportion of CO₂ emissions taken up by land and ocean carbon sinks is smaller in scenarios with higher cumulative CO₂ emissions

Attachment 1

Total cumulative CO₂ emissions taken up by land and oceans (colours) and remaining in the atmosphere (grey) under the five illustrative scenarios from 1850 to 2100




Figure SPM.7: Cumulative anthropogenic CO₂ emissions taken up by land and ocean sinks by 2100 under the five illustrative scenarios.

The cumulative anthropogenic (human-caused) carbon dioxide (CO₂) emissions taken up by the land and ocean sinks under the five illustrative scenarios (SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP3-7.0 and SSP5-8.5) are simulated from 1850 to 2100 by CMIP6 climate models in the concentration-driven simulations. Land and ocean carbon sinks respond to past, current and future emissions, therefore cumulative sinks from 1850 to 2100 are presented here. During the historical period (1850-2019) the observed land and ocean sink took up 1430 GtCO₂ (59% of the emissions).

The **bar chart** illustrates the projected amount of cumulative anthropogenic CO₂ emissions (GtCO₂) between 1850 and 2100 remaining in the atmosphere (grey part) and taken up by the land and ocean (coloured part) in the year 2100. The **doughnut chart** illustrates the proportion of the cumulative anthropogenic CO₂ emissions taken up by the land and ocean sinks and remaining in the atmosphere in the year 2100. Values in % indicate the proportion of the cumulative anthropogenic CO₂ emissions taken up by the combined land and ocean sinks in the year 2100. The overall anthropogenic carbon emissions are calculated by adding the net global land use emissions from CMIP6 scenario database to the other sectoral emissions calculated from climate model runs with prescribed CO₂ concentrations³³. Land and ocean CO₂ uptake since 1850 is calculated from the net biome productivity on land, corrected for CO₂ losses due to land-use change by adding the land-use change emissions, and net ocean CO₂ flux.

{Box TS.5, Box TS.5, Figure 1, 5.2.1, Table 5.1, 5.4.5, Figure 5.25}

³³ The other sectoral emissions are calculated as the residual of the net land and ocean CO₂ uptake and the prescribed atmospheric CO₂ concentration changes in the CMIP6 simulations. These calculated emissions are net emissions and do not separate gross anthropogenic emissions from removals, which are included implicitly.

B.5 Many changes due to past and future greenhouse gas emissions are irreversible for centuries to millennia, especially changes in the ocean, ice sheets and global sea level.

{Cross-Chapter Box 2.4, 2.3, 4.3, 4.5, 4.7, 5.3, 9.2, 9.4, 9.5, 9.6, Box 9.4} (Figure SPM.8)

Attachment 1

B.5.1 Past GHG emissions since 1750 have committed the global ocean to future warming (*high confidence*). Over the rest of the 21st century, *likely* ocean warming ranges from 2–4 (SSP1-2.6) to 4–8 times (SSP5-8.5) the 1971–2018 change. Based on multiple lines of evidence, upper ocean stratification (*virtually certain*), ocean acidification (*virtually certain*) and ocean deoxygenation (*high confidence*) will continue to increase in the 21st century, at rates dependent on future emissions. Changes are irreversible on centennial to millennial time scales in global ocean temperature (*very high confidence*), deep ocean acidification (*very high confidence*) and deoxygenation (*medium confidence*). {4.3, 4.5, 4.7, 5.3, 9.2, TS.2.4} (Figure SPM.8)

B.5.2 Mountain and polar glaciers are committed to continue melting for decades or centuries (*very high confidence*). Loss of permafrost carbon following permafrost thaw is irreversible at centennial timescales (*high confidence*). Continued ice loss over the 21st century is *virtually certain* for the Greenland Ice Sheet and *likely* for the Antarctic Ice Sheet. There is *high confidence* that total ice loss from the Greenland Ice Sheet will increase with cumulative emissions. There is *limited evidence* for low-likelihood, high-impact outcomes (resulting from ice sheet instability processes characterized by deep uncertainty and in some cases involving tipping points) that would strongly increase ice loss from the Antarctic Ice Sheet for centuries under high GHG emissions scenarios³⁴. {4.3, 4.7, 5.4, 9.4, 9.5, Box 9.4, Box TS.1, TS.2.5}

B.5.3 It is *virtually certain* that global mean sea level will continue to rise over the 21st century. Relative to 1995–2014, the *likely* global mean sea level rise by 2100 is 0.28–0.55 m under the very low GHG emissions scenario (SSP1-1.9), 0.32–0.62 m under the low GHG emissions scenario (SSP1-2.6), 0.44–0.76 m under the intermediate GHG emissions scenario (SSP2-4.5), and 0.63–1.01 m under the very high GHG emissions scenario (SSP5-8.5), and by 2150 is 0.37–0.86 m under the very low scenario (SSP1-1.9), 0.46–0.99 m under the low scenario (SSP1-2.6), 0.66–1.33 m under the intermediate scenario (SSP2-4.5), and 0.98–1.88 m under the very high scenario (SSP5-8.5) (*medium confidence*)³⁵. Global mean sea level rise above the *likely* range – approaching 2 m by 2100 and 5 m by 2150 under a very high GHG emissions scenario (SSP5-8.5) (*low confidence*) – cannot be ruled out due to deep uncertainty in ice sheet processes. {4.3, 9.6, Box 9.4, Box TS.4} (Figure SPM.8)

B.5.4 In the longer term, sea level is committed to rise for centuries to millennia due to continuing deep ocean warming and ice sheet melt, and will remain elevated for thousands of years (*high confidence*). Over the next 2000 years, global mean sea level will rise by about 2 to 3 m if warming is limited to 1.5°C, 2 to 6 m if limited to 2°C and 19 to 22 m with 5°C of warming, and it will continue to rise over subsequent millennia (*low confidence*). Projections of multi-millennial global mean sea level rise are consistent with reconstructed levels during past warm climate periods: *likely* 5–10 m higher than today around 125,000 years ago, when global temperatures were *very likely* 0.5°C–1.5°C higher than 1850–1900; and *very likely* 5–25 m higher roughly 3 million years ago, when global temperatures were 2.5°C–4°C higher (*medium confidence*). {2.3, Cross-Chapter Box 2.4, 9.6, Box TS.2, Box TS.4, Box TS.9}


³⁴ Low-likelihood, high-impact outcomes are those whose probability of occurrence is low or not well known (as in the context of deep uncertainty) but whose potential impacts on society and ecosystems could be high. A tipping point is a critical threshold beyond which a system reorganizes, often abruptly and/or irreversibly. {Cross-Chapter Box 1.3, 1.4, 4.7}

³⁵ To compare to the 1986–2005 baseline period used in AR5 and SROCC, add 0.03 m to the global mean sea level rise estimates. To compare to the 1900 baseline period used in Figure SPM.8, add 0.16 m.


Human activities affect all the major climate system components, with some responding over decades and others over centuries

Attachment 1


a) Global surface temperature change relative to 1850-1900




b) September Arctic sea ice area



c) Global ocean surface pH (a measure of acidity)



d) Global mean sea level change relative to 1900



e) Global mean sea level change in 2300 relative to 1900




Figure SPM.8: Selected indicators of global climate change under the five illustrative scenarios used in this report. Attachment 1

The projections for each of the five scenarios are shown in colour. Shades represent uncertainty ranges – more detail is provided for each panel below. The black curves represent the historical simulations (panels a, b, c) or the observations (panel d). Historical values are included in all graphs to provide context for the projected future changes.

Panel a) Global surface temperature changes in °C relative to 1850–1900. These changes were obtained by combining CMIP6 model simulations with observational constraints based on past simulated warming, as well as an updated assessment of equilibrium climate sensitivity (see Box SPM.1). Changes relative to 1850–1900 based on 20-year averaging periods are calculated by adding 0.85°C (the observed global surface temperature increase from 1850–1900 to 1995–2014) to simulated changes relative to 1995–2014. *Very likely* ranges are shown for SSP1-2.6 and SSP3-7.0.

Panel b) September Arctic sea ice area in 10⁶ km² based on CMIP6 model simulations. *Very likely* ranges are shown for SSP1-2.6 and SSP3-7.0. The Arctic is projected to be practically ice-free near mid-century under mid- and high GHG emissions scenarios.

Panel c) Global ocean surface pH (a measure of acidity) based on CMIP6 model simulations. *Very likely* ranges are shown for SSP1-2.6 and SSP3-7.0.

Panel d) Global mean sea level change in meters relative to 1900. The historical changes are observed (from tide gauges before 1992 and altimeters afterwards), and the future changes are assessed consistently with observational constraints based on emulation of CMIP, ice sheet, and glacier models. *Likely* ranges are shown for SSP1-2.6 and SSP3-7.0. Only *likely* ranges are assessed for sea level changes due to difficulties in estimating the distribution of deeply uncertain processes. The dashed curve indicates the potential impact of these deeply uncertain processes. It shows the 83rd percentile of SSP5-8.5 projections that include low-likelihood, high-impact ice sheet processes that cannot be ruled out; because of *low confidence* in projections of these processes, this curve does not constitute part of a *likely* range. Changes relative to 1900 are calculated by adding 0.158 m (observed global mean sea level rise from 1900 to 1995–2014) to simulated and observed changes relative to 1995–2014.

Panel e): Global mean sea level change at 2300 in meters relative to 1900. Only SSP1-2.6 and SSP5-8.5 are projected at 2300, as simulations that extend beyond 2100 for the other scenarios are too few for robust results. The 17th–83rd percentile ranges are shaded. The dashed arrow illustrates the 83rd percentile of SSP5-8.5 projections that include low-likelihood, high-impact ice sheet processes that cannot be ruled out.

Panels b) and c) are based on single simulations from each model, and so include a component of internal variability. Panels a), d) and e) are based on long-term averages, and hence the contributions from internal variability are small.

{Figure TS.8, Figure TS.11, Box TS.4 Figure 1, Box TS.4 Figure 1, 4.3, 9.6, Figure 4.2, Figure 4.8, Figure 4.11, Figure 9.27}

C. Climate Information for Risk Assessment and Regional Adaptation

Attachment 1

Physical climate information addresses how the climate system responds to the interplay between human influence, natural drivers and internal variability. Knowledge of the climate response and the range of possible outcomes, including low-likelihood, high impact outcomes, informs climate services – the assessment of climate-related risks and adaptation planning. Physical climate information at global, regional and local scales is developed from multiple lines of evidence, including observational products, climate model outputs and tailored diagnostics.

C.1 Natural drivers and internal variability will modulate human-caused changes, especially at regional scales and in the near term, with little effect on centennial global warming. These modulations are important to consider in planning for the full range of possible changes.

{1.4, 2.2, 3.3, Cross-Chapter Box 3.1, 4.4, 4.6, Cross-Chapter Box 4.1, 4.4, Box 7.2, 8.3, 8.5, 9.2, 10.3, 10.4, 10.6, 11.3, 12.5, Atlas.4, Atlas.5, Atlas.8, Atlas.9, Atlas.10, Cross-Chapter Box Atlas.2, Atlas.11}

C.1.1 The historical global surface temperature record highlights that decadal variability has enhanced and masked underlying human-caused long-term changes, and this variability will continue into the future (*very high confidence*). For example, internal decadal variability and variations in solar and volcanic drivers partially masked human-caused surface global warming during 1998–2012, with pronounced regional and seasonal signatures (*high confidence*). Nonetheless, the heating of the climate system continued during this period, as reflected in both the continued warming of the global ocean (*very high confidence*) and in the continued rise of hot extremes over land (*medium confidence*).

{1.4, 3.3, Cross-Chapter Box 3.1, 4.4, Box 7.2, 9.2, 11.3, Cross-Section Box TS.1} (**Figure SPM.1**)

C.1.2 Projected human caused changes in mean climate and climatic impact-drivers (CIDs)³⁶, including extremes, will be either amplified or attenuated by internal variability³⁷ (*high confidence*). Near-term cooling at any particular location with respect to present climate could occur and would be consistent with the global surface temperature increase due to human influence (*high confidence*).

{1.4, 4.4, 4.6, 10.4, 11.3, 12.5, Atlas.5, Atlas.10, Atlas.11, TS.4.2}

C.1.3 Internal variability has largely been responsible for the amplification and attenuation of the observed human-caused decadal-to-multi-decadal mean precipitation changes in many land regions (*high confidence*). At global and regional scales, near-term changes in monsoons will be dominated by the effects of internal variability (*medium confidence*). In addition to internal variability influence, near-term projected changes in precipitation at global and regional scales are uncertain because of model uncertainty and uncertainty in forcings from natural and anthropogenic aerosols (*medium confidence*).

{1.4, 4.4, 8.3, 8.5, 10.3, 10.4, 10.5, 10.6, Atlas.4, Atlas.8, Atlas.9, Atlas.10, Cross-Chapter Box Atlas.2, Atlas.11, TS.4.2, Box TS.6, Box TS.13}

³⁶ Climatic impact-drivers (CIDs) are physical climate system conditions (e.g., means, events, extremes) that affect an element of society or ecosystems. Depending on system tolerance, CIDs and their changes can be detrimental, beneficial, neutral, or a mixture of each across interacting system elements and regions. CID types include heat and cold, wet and dry, wind, snow and ice, coastal and open ocean.

³⁷ The main internal variability phenomena include El Niño–Southern Oscillation, Pacific Decadal variability and Atlantic Multi-decadal variability through their regional influence.

C.1.4 Based on paleoclimate and historical evidence, it is *likely* that at least one large explosive volcanic eruption would occur during the 21st century³⁸. Such an eruption would reduce global surface temperature and precipitation, especially over land, for one to three years, alter the global monsoon circulation, modify extreme precipitation and change many CIDs (*medium confidence*). If such an eruption occurs, this would therefore temporarily and partially mask human-caused climate change. Attachment 1
{4.4, Cross-Chapter Box 4.1, 2.2, 8.5, TS.2.1}

C.2 With further global warming, every region is projected to increasingly experience concurrent and multiple changes in climatic impact-drivers. Changes in several climatic impact-drivers would be more widespread at 2°C compared to 1.5°C global warming and even more widespread and/or pronounced for higher warming levels.

{8.2, 9.3, 9.5, 9.6, Box 10.3, Box 11.3, Box 11.4, 11.3, 11.4, 11.5, 11.6, 11.7, 11.9, 12.2, 12.3, 12.4, 12.5, Atlas.4, Atlas.5, Atlas.6, Atlas.7, Atlas.8, Atlas.9, Atlas.10, Atlas.11, Cross-Chapter Box 11.1, Cross-Chapter Box 12.1} (Table SPM.1, Figure SPM.9)

C.2.1 All regions³⁹ are projected to experience further increases in hot climatic impact-drivers (CIDs) and decreases in cold CIDs (*high confidence*). Further decreases are projected in permafrost, snow, glaciers and ice sheets, lake and Arctic sea ice (*medium to high confidence*)⁴⁰. These changes would be larger at 2°C global warming or above than at 1.5°C (*high confidence*). For example, extreme heat thresholds relevant to agriculture and health are projected to be exceeded more frequently at higher global warming levels (*high confidence*).

{9.3, 9.5, 11.3, 11.9, 12.3, 12.4, 12.5, Atlas.4, Atlas.5, Atlas.6, Atlas.7, Atlas.8, Atlas.9, Atlas.10, Atlas.11, TS.4.3, Cross-Chapter Box 11.1, Cross-Chapter Box 12.1} (Table SPM.1, Figure SPM.9)

C.2.2 At 1.5°C global warming, heavy precipitation and associated flooding are projected to intensify and be more frequent in most regions in Africa and Asia (*high confidence*), North America (*medium to high confidence*)⁴⁰ and Europe (*medium confidence*). Also, more frequent and/or severe agricultural and ecological droughts are projected in a few regions in all continents except Asia compared to 1850–1900 (*medium confidence*); increases in meteorological droughts are also projected in a few regions (*medium confidence*). A small number of regions are projected to experience increases or decreases in mean precipitation (*medium confidence*).

{11.4, 11.5, 11.6, 11.9, Atlas.4, Atlas.5, Atlas.7, Atlas.8, Atlas.9, Atlas.10, Atlas.11, TS.4.3} (Table SPM.1)

³⁸ Based on 2,500 year reconstructions, eruptions more negative than -1 W m^{-2} occur on average twice per century.

³⁹ Regions here refer to the AR6 WGI reference regions used in this Report to summarize information in sub-continental and oceanic regions. Changes are compared to averages over the last 20–40 years unless otherwise specified. {1.4, 12.4, Atlas.1, Interactive Atlas}.

⁴⁰ The specific level of confidence or likelihood depends on the region considered. Details can be found in the Technical Summary and the underlying Report.

C.2.3 At 2°C global warming and above, the level of confidence in and the magnitude of the change in droughts and heavy and mean precipitation increase compared to those at 1.5°C. Heavy precipitation and associated flooding events are projected to become more intense and frequent in the Pacific Islands and across many regions of North America and Europe (*medium to high confidence*)⁴⁰. These changes are also seen in some regions in Australasia and Central and South America (*medium confidence*). Several regions in Africa, South America and Europe are projected to experience an increase in frequency and/or severity of agricultural and ecological droughts with *medium to high confidence*⁴⁰; increases are also projected in Australasia, Central and North America, and the Caribbean with *medium confidence*. A small number of regions in Africa, Australasia, Europe and North America are also projected to be affected by increases in hydrological droughts, and several regions are projected to be affected by increases or decreases in meteorological droughts with more regions displaying an increase (*medium confidence*). Mean precipitation is projected to increase in all polar, northern European and northern North American regions, most Asian regions and two regions of South America (*high confidence*).
{11.4, 11.6, 11.9, 12.4, 12.5, Atlas.5, Atlas.7, Atlas.8, Atlas.9, Atlas.11, TS.4.3, Cross-Chapter Box 11.1, Cross-Chapter Box 12.1} (**Table SPM.1, Figure SPM.5, Figure SPM.6, Figure SPM.9**)

C.2.4 More CIDs across more regions are projected to change at 2°C and above compared to 1.5°C global warming (*high confidence*). Region-specific changes include intensification of tropical cyclones and/or extratropical storms (*medium confidence*), increases in river floods (*medium to high confidence*)⁴⁰, reductions in mean precipitation and increases in aridity (*medium to high confidence*)⁴⁰, and increases in fire weather (*medium to high confidence*)⁴⁰. There is *low confidence* in most regions in potential future changes in other CIDs, such as hail, ice storms, severe storms, dust storms, heavy snowfall, and landslides.
{11.7, 11.9, 12.4, 12.5, Atlas.4, Atlas.6, Atlas.7, Atlas.8, Atlas.10, TS.4.3.1, TS.4.3.2, TS.5, Cross-Chapter Box, 11.1, Cross-Chapter Box 12.1} (**Table SPM.1, Figure SPM.9**)

C.2.5 It is *very likely to virtually certain*⁴⁰ that regional mean relative sea level rise will continue throughout the 21st century, except in a few regions with substantial geologic land uplift rates. Approximately two-thirds of the global coastline has a projected regional relative sea level rise within ±20% of the global mean increase (*medium confidence*). Due to relative sea level rise, extreme sea level events that occurred once per century in the recent past are projected to occur at least annually at more than half of all tide gauge locations by 2100 (*high confidence*). Relative sea level rise contributes to increases in the frequency and severity of coastal flooding in low-lying areas and to coastal erosion along most sandy coasts (*high confidence*).
{9.6, 12.4, 12.5, Box TS.4, TS.4.3, Cross-Chapter Box 12.1} (**Figure SPM.9**)


C.2.6 Cities intensify human-induced warming locally, and further urbanization together with more frequent hot extremes will increase the severity of heatwaves (*very high confidence*). Urbanization also increases mean and heavy precipitation over and/or downwind of cities (*medium confidence*) and resulting runoff intensity (*high confidence*). In coastal cities, the combination of more frequent extreme sea level events (due to sea level rise and storm surge) and extreme rainfall/riverflow events will make flooding more probable (*high confidence*).
{8.2, Box 10.3, 11.3, 12.4, Box TS.14}

C.2.7 Many regions are projected to experience an increase in the probability of compound events with higher global warming (*high confidence*). In particular, concurrent heatwaves and droughts are *likely* to become more frequent. Concurrent extremes at multiple locations become more frequent, including in crop-producing areas, at 2°C and above compared to 1.5°C global warming (*high confidence*).
{11.8, Box 11.3, Box 11.4, 12.3, 12.4, TS.4.3, Cross-Chapter Box 12.1} (**Table SPM.1**)

Multiple climatic impact-drivers are projected to change in all regions of the world


Attachment 1

Climatic impact-drivers (CIDs) are physical climate system conditions (e.g., means, events, extremes) that affect an element of society or ecosystems. Depending on system tolerance, CIDs and their changes can be detrimental, beneficial, neutral, or a mixture of each across interacting system elements and regions. The CIDs are grouped into seven types, which are summarized under the icons in the figure. All regions are projected to experience changes in at least 5 CIDs. Almost all (96%) are projected to experience changes in at least 10 CIDs and half in at least 15 CIDs. For many CIDs there is wide geographical variation in where they change and so each region are projected to experience a specific set of CID changes. Each bar in the chart represents a specific geographical set of changes that can be explored in the WGI Interactive Atlas.



interactive-atlas.ipcc.ch

Number of land & coastal regions (a) and open-ocean regions (b) where each climatic impact-driver (CID) is projected to increase or decrease with high confidence (dark shade) or medium confidence (light shade)



BAR CHART LEGEND

- Regions with **high confidence increase**
- Regions with **medium confidence increase**
- Regions with **high confidence decrease**
- Regions with **medium confidence decrease**

LIGHTER-SHADED 'ENVELOPE' LEGEND

The height of the lighter shaded 'envelope' behind each bar represents the maximum number of regions for which each CID is relevant. The envelope is symmetrical about the x-axis showing the maximum possible number of relevant regions for CID increase (upper part) or decrease (lower part).

ASSESSED FUTURE CHANGES

Changes refer to a 20–30 year period centred around 2050 and/or consistent with 2°C global warming compared to a similar period within 1960–2014 or 1850–1900.

Figure SPM.9: Synthesis of the number of AR6 WGI reference regions where climatic impact-drivers are projected to change.

Attachment 1

A total of 35 climatic impact-drivers (CIDs) grouped into seven types are shown: heat and cold, wet and dry, wind, snow and ice, coastal, open ocean and other. For each CID, the bar in the graph below displays the number of AR6 WGI reference regions where it is projected to change. The **colours** represent the direction of change and the level of confidence in the change: purple indicates an increase while brown indicates a decrease; darker and lighter shades refer to *high* and *medium confidence*, respectively. Lighter background colours represent the maximum number of regions for which each CID is broadly relevant.

Panel a) shows the 30 CIDs relevant to the **land and coastal regions** while **panel b)** shows the 5 CIDs relevant to the **open ocean regions**. Marine heatwaves and ocean acidity are assessed for coastal ocean regions in panel a) and for open ocean regions in panel b). Changes refer to a 20–30 year period centred around 2050 and/or consistent with 2°C global warming compared to a similar period within 1960-2014, except for hydrological drought and agricultural and ecological drought which is compared to 1850-1900. Definitions of the regions are provided in Atlas.1 and the Interactive Atlas (see *interactive-atlas.ipcc.ch*).

{Table TS.5, Figure TS.22, Figure TS.25, 11.9, 12.2, 12.4, Atlas.1} **(Table SPM.1)**

C.3 Low-likelihood outcomes, such as ice sheet collapse, abrupt ocean circulation changes, some compound extreme events and warming substantially larger than the assessed *very likely* range of future warming cannot be ruled out and are part of risk assessment.

{1.4, Cross-Chapter Box 1.3, Cross-Chapter Box 4.1, 4.3, 4.4, 4.8, 8.6, 9.2, Box 9.4, Box 11.2, 11.8, Cross-Chapter Box 12.1} **(Table SPM.1)**

C.3.1 If global warming exceeds the assessed *very likely* range for a given GHG emissions scenario, including low GHG emissions scenarios, global and regional changes in many aspects of the climate system, such as regional precipitation and other CIDs, would also exceed their assessed *very likely* ranges (*high confidence*). Such low-likelihood high-warming outcomes are associated with potentially very large impacts, such as through more intense and more frequent heatwaves and heavy precipitation, and high risks for human and ecological systems particularly for high GHG emissions scenarios.

{Cross-Chapter Box 1.3, 4.3, 4.4, 4.8, Box 9.4, Box 11.2, Cross-Chapter Box 12.1, TS.1.4, Box TS.3, Box TS.4} **(Table SPM.1)**

C.3.2 Low-likelihood, high-impact outcomes³⁴ could occur at global and regional scales even for global warming within the *very likely* range for a given GHG emissions scenario. The probability of low-likelihood, high impact outcomes increases with higher global warming levels (*high confidence*). Abrupt responses and tipping points of the climate system, such as strongly increased Antarctic ice sheet melt and forest dieback, cannot be ruled out (*high confidence*).

{1.4, 4.3, 4.4, 4.8, 5.4, 8.6, Box 9.4, Cross-Chapter Box 12.1, TS.1.4, TS.2.5, Box TS.3, Box TS.4, Box TS.9} **(Table SPM.1)**

C.3.3 If global warming increases, some compound extreme events¹⁸ with low likelihood in past and current climate will become more frequent, and there will be a higher likelihood that events with increased intensities, durations and/or spatial extents unprecedented in the observational record will occur (*high confidence*).

{11.8, Box 11.2, Cross-Chapter Box 12.1, Box TS.3, Box TS.9}

C.3.4 The Atlantic Meridional Overturning Circulation is *very likely* to weaken over the 21st century for all emission scenarios. While there is *high confidence* in the 21st century decline, there is only *low confidence* in the magnitude of the trend. There is *medium confidence* that there will not be an abrupt collapse before 2100. If such a collapse were to occur, it would *very likely* cause abrupt shifts in regional weather patterns and water cycle, such as a southward shift in the tropical rain belt, weakening of the African and Asian monsoons and strengthening of Southern Hemisphere monsoons, and drying in Europe. {4.3, 8.6, 9.2, TS2.4, Box TS.3}

C.3.5 Unpredictable and rare natural events not related to human influence on climate may lead to low-likelihood, high impact outcomes. For example, a sequence of large explosive volcanic eruptions within decades has occurred in the past, causing substantial global and regional climate perturbations over several decades. Such events cannot be ruled out in the future, but due to their inherent unpredictability they are not included in the illustrative set of scenarios referred to in this Report. {2.2, Cross-Chapter Box 4.1, Box TS.3} **(Box SPM.1)**

D. Limiting Future Climate Change

Since AR5, estimates of remaining carbon budgets have been improved by a new methodology first presented in SR1.5, updated evidence, and the integration of results from multiple lines of evidence. A comprehensive range of possible future air pollution controls in scenarios is used to consistently assess the effects of various assumptions on projections of climate and air pollution. A novel development is the ability to ascertain when climate responses to emissions reductions would become discernible above natural climate variability, including internal variability and responses to natural drivers.

D.1 From a physical science perspective, limiting human-induced global warming to a specific level requires limiting cumulative CO₂ emissions, reaching at least net zero CO₂ emissions, along with strong reductions in other greenhouse gas emissions. Strong, rapid and sustained reductions in CH₄ emissions would also limit the warming effect resulting from declining aerosol pollution and would improve air quality. {3.3, 4.6, 5.1, 5.2, 5.4, 5.5, 5.6, Box 5.2, Cross-Chapter Box 5.1, 6.7, 7.6, 9.6} **(Figure SPM.10, Table SPM.2)**

D.1.1 This Report reaffirms with *high confidence* the AR5 finding that there is a near-linear relationship between cumulative anthropogenic CO₂ emissions and the global warming they cause. Each 1000 GtCO₂ of cumulative CO₂ emissions is assessed to *likely* cause a 0.27°C to 0.63°C increase in global surface temperature with a best estimate of 0.45°C⁴¹. This is a narrower range compared to AR5 and SR1.5. This quantity is referred to as the transient climate response to cumulative CO₂ emissions (TCRE). This relationship implies that reaching net zero⁴² anthropogenic CO₂ emissions is a requirement to stabilize human-induced global temperature increase at any level, but that limiting global temperature increase to a specific level would imply limiting cumulative CO₂ emissions to within a carbon budget⁴³. {5.4, 5.5, TS.1.3, TS.3.3, Box TS.5} **(Figure SPM.10)**

⁴¹ In the literature, units of °C per 1000 PgC are used, and the AR6 reports the TCRE *likely* range as 1.0°C to 2.3°C per 1000 PgC in the underlying report, with a best estimate of 1.65°C.

⁴² condition in which anthropogenic carbon dioxide (CO₂) emissions are balanced by anthropogenic CO₂ removals over a specified period.

⁴³ The term carbon budget refers to the maximum amount of cumulative net global anthropogenic CO₂ emissions that would result in limiting global warming to a given level with a given probability, taking into account the effect of other anthropogenic climate forcings. This is referred to as the total carbon budget when expressed starting from the pre-industrial period, and as the remaining carbon budget when expressed from a recent specified date (see Glossary). Historical cumulative CO₂ emissions determine to a large degree warming to date, while future emissions cause future additional warming. The remaining carbon budget indicates how much CO₂ could still be emitted while keeping warming below a specific temperature level.

Every tonne of CO₂ emissions adds to global warming

Global surface temperature increase since 1850-1900 (°C) as a function of cumulative CO₂ emissions (GtCO₂) Attachment 1




Figure SPM.10: Near-linear relationship between cumulative CO₂ emissions and the increase in global surface temperature.

Top panel: Historical data (thin black line) shows observed global surface temperature increase in °C since 1850–1900 as a function of historical cumulative carbon dioxide (CO₂) emissions in GtCO₂ from 1850 to 2019. The grey range with its central line shows a corresponding estimate of the historical human-caused surface warming (see Figure SPM.2). Coloured areas show the assessed *very likely* range of global surface temperature projections, and thick coloured central lines show the median estimate as a function of cumulative CO₂ emissions from 2020 until year 2050 for the set of illustrative scenarios (SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5, see Figure SPM.4). Projections use the cumulative CO₂ emissions of each respective scenario, and the projected global warming includes the contribution from all anthropogenic forcings. The relationship is illustrated over the domain of cumulative CO₂ emissions for which there is *high confidence* that the transient climate response to cumulative CO₂ emissions (TCRE) remains constant, and for the time period from 1850 to 2050 over which global CO₂ emissions remain net positive under all illustrative scenarios as there is *limited evidence* supporting the quantitative application of TCRE to estimate temperature evolution under net negative CO₂ emissions.

Bottom panel: Historical and projected cumulative CO₂ emissions in GtCO₂ for the respective scenarios.

{Figure TS.18, Figure 5.31, Section 5.5}

D.1.2 Over the period 1850–2019, a total of 2390 ± 240 (*likely* range) GtCO₂ of anthropogenic CO₂ was emitted. Remaining carbon budgets have been estimated for several global temperature limits and various levels of probability, based on the estimated value of TCRE and its uncertainty, estimates of historical warming, variations in projected warming from non-CO₂ emissions, climate system feedbacks such as emissions from thawing permafrost, and the global surface temperature change after global anthropogenic CO₂ emissions reach net zero. {5.1, 5.5, Box 5.2, TS.3.3} (**Table SPM.2**)

Table SPM.2: Estimates of historical CO₂ emissions and remaining carbon budgets. Estimated remaining carbon budgets are calculated from the beginning of 2020 and extend until global net zero CO₂ emissions are reached. They refer to CO₂ emissions, while accounting for the global warming effect of non-CO₂ emissions. Global warming in this table refers to human-induced global surface temperature increase, which excludes the impact of natural variability on global temperatures in individual years. {Table TS.3, Table 3.1, Table 5.1, Table 5.7, Table 5.8, 5.5.1, 5.5.2, Box 5.2}

Global warming between 1850–1900 and 2010–2019 (°C)	Historical cumulative CO ₂ emissions from 1850 to 2019 (GtCO ₂)
1.07 (0.8–1.3; <i>likely</i> range)	2390 (± 240; <i>likely</i> range)

Approximate global warming relative to 1850–1900 until temperature limit (°C)* ⁽¹⁾	Additional global warming relative to 2010–2019 until temperature limit (°C)	Estimated remaining carbon budgets from the beginning of 2020 (GtCO ₂)					Variations in reductions in non-CO ₂ emissions* ⁽³⁾
		<i>Likelihood of limiting global warming to temperature limit*⁽²⁾</i>					
		17%	33%	50%	67%	83%	
1.5	0.43	900	650	500	400	300	Higher or lower reductions in accompanying non-CO ₂ emissions can increase or decrease the values on the left by 220 GtCO ₂ or more
1.7	0.63	1450	1050	850	700	550	
2.0	0.93	2300	1700	1350	1150	900	

*⁽¹⁾ Values at each 0.1°C increment of warming are available in Tables TS.3 and 5.8.
 *⁽²⁾ This likelihood is based on the uncertainty in transient climate response to cumulative CO₂ emissions (TCRE) and additional Earth system feedbacks, and provides the probability that global warming will not exceed the temperature levels provided in the two left columns. Uncertainties related to historical warming (±550 GtCO₂) and non-CO₂ forcing and response (±220 GtCO₂) are partially addressed by the assessed uncertainty in TCRE, but uncertainties in recent emissions since 2015 (±20 GtCO₂) and the climate response after net zero CO₂ emissions are reached (±420 GtCO₂) are separate.
 *⁽³⁾ Remaining carbon budget estimates consider the warming from non-CO₂ drivers as implied by the scenarios assessed in SR1.5. The Working Group III Contribution to AR6 will assess mitigation of non-CO₂ emissions.

D.1.3 Several factors that determine estimates of the remaining carbon budget have been re-assessed, and updates to these factors since SR1.5 are small. When adjusted for emissions since previous reports, estimates of remaining carbon budgets are therefore of similar magnitude compared to SR1.5 but larger compared to AR5 due to methodological improvements⁴⁴. Attachment 1
{5.5, Box 5.2, TS.3.3} (**Table SPM.2**)

D.1.4 Anthropogenic CO₂ removal (CDR) has the potential to remove CO₂ from the atmosphere and durably store it in reservoirs (*high confidence*). CDR aims to compensate for residual emissions to reach net zero CO₂ or net zero GHG emissions or, if implemented at a scale where anthropogenic removals exceed anthropogenic emissions, to lower surface temperature. CDR methods can have potentially wide-ranging effects on biogeochemical cycles and climate, which can either weaken or strengthen the potential of these methods to remove CO₂ and reduce warming, and can also influence water availability and quality, food production and biodiversity⁴⁵ (*high confidence*).
{5.6, Cross-Chapter Box 5.1, TS.3.3}

D.1.5 Anthropogenic CO₂ removal (CDR) leading to global net negative emissions would lower the atmospheric CO₂ concentration and reverse surface ocean acidification (*high confidence*). Anthropogenic CO₂ removals and emissions are partially compensated by CO₂ release and uptake respectively, from or to land and ocean carbon pools (*very high confidence*). CDR would lower atmospheric CO₂ by an amount approximately equal to the increase from an anthropogenic emission of the same magnitude (*high confidence*). The atmospheric CO₂ decrease from anthropogenic CO₂ removals could be up to 10% less than the atmospheric CO₂ increase from an equal amount of CO₂ emissions, depending on the total amount of CDR (*medium confidence*). {5.3, 5.6, TS.3.3}

D.1.6 If global net negative CO₂ emissions were to be achieved and be sustained, the global CO₂-induced surface temperature increase would be gradually reversed but other climate changes would continue in their current direction for decades to millennia (*high confidence*). For instance, it would take several centuries to millennia for global mean sea level to reverse course even under large net negative CO₂ emissions (*high confidence*).
{4.6, 9.6, TS.3.3}

D.1.7 In the five illustrative scenarios, simultaneous changes in CH₄, aerosol and ozone precursor emissions, that also contribute to air pollution, lead to a net global surface warming in the near and long-term (*high confidence*). In the long term, this net warming is lower in scenarios assuming air pollution controls combined with strong and sustained CH₄ emission reductions (*high confidence*). In the low and very low GHG emissions scenarios, assumed reductions in anthropogenic aerosol emissions lead to a net warming, while reductions in CH₄ and other ozone precursor emissions lead to a net cooling. Because of the short lifetime of both CH₄ and aerosols, these climate effects partially counterbalance each other and reductions in CH₄ emissions also contribute to improved air quality by reducing global surface ozone (*high confidence*).
{6.7, Box TS.7} (**Figure SPM.2, Box SPM.1**)

⁴⁴ Compared to AR5, and when taking into account emissions since AR5, estimates in AR6 are about 300–350 GtCO₂ larger for the remaining carbon budget consistent with limiting warming to 1.5°C; for 2°C, the difference is about 400–500 GtCO₂.

⁴⁵ Potential negative and positive effects of CDR for biodiversity, water and food production are methods-specific, and are often highly dependent on local context, management, prior land use, and scale. IPCC Working Groups II and III assess the CDR potential, and ecological and socio-economic effects of CDR methods in their AR6 contributions.

D.1.8 Achieving global net zero CO₂ emissions is a requirement for stabilizing CO₂-induced global surface temperature increase, with anthropogenic CO₂ emissions balanced by anthropogenic removals of CO₂. This is different from achieving net zero GHG emissions, where metric-weighted anthropogenic GHG emissions are balanced by equal metric-weighted anthropogenic GHG removals. For a given GHG emission pathway, the pathways of individual greenhouse gases determine the resulting climate response⁴⁶, whereas the choice of emissions metric⁴⁷ used to calculate aggregated emissions and removals of different GHGs affects what point in time the aggregated greenhouse gases are calculated to be net zero. Emissions pathways that reach and sustain net zero GHG emissions defined by the 100-year global warming potential are projected to result in a decline in surface temperature after an earlier peak (*high confidence*).
{4.6, 7.6, Box 7.3, TS.3.3}

D.2 Scenarios with very low or low GHG emissions (SSP1-1.9 and SSP1-2.6) lead within years to discernible effects on greenhouse gas and aerosol concentrations, and air quality, relative to high and very high GHG emissions scenarios (SSP3-7.0 or SSP5-8.5). Under these contrasting scenarios, discernible differences in trends of global surface temperature would begin to emerge from natural variability within around 20 years, and over longer time periods for many other climatic impact-drivers (*high confidence*).
{4.6, Cross-Chapter Box 6.1, 6.6, 6.7, 9.6, Cross-Chapter Box 11.1, 11.2, 11.4, 11.5, 11.6, 12.4, 12.5} (Figure SPM.8, Figure SPM.10)

D.2.1 Emissions reductions in 2020 associated with measures to reduce the spread of COVID-19 led to temporary but detectable effects on air pollution (*high confidence*), and an associated small, temporary increase in total radiative forcing, primarily due to reductions in cooling caused by aerosols arising from human activities (*medium confidence*). Global and regional climate responses to this temporary forcing are, however, undetectable above natural variability (*high confidence*). Atmospheric CO₂ concentrations continued to rise in 2020, with no detectable decrease in the observed CO₂ growth rate (*medium confidence*)⁴⁸.
{Cross-Chapter Box 6.1, TS.3.3}

D.2.2 Reductions in GHG emissions also lead to air quality improvements. However, in the near term⁴⁹, even in scenarios with strong reduction of GHGs, as in the low and very low GHG emission scenarios (SSP1-2.6 and SSP1-1.9), these improvements are not sufficient in many polluted regions to achieve air quality guidelines specified by the World Health Organization (*high confidence*). Scenarios with targeted reductions of air pollutant emissions lead to more rapid improvements in air quality within years compared to reductions in GHG emissions only, but from 2040, further improvements are projected in scenarios that combine efforts to reduce air pollutants as well as GHG emissions with the magnitude of the benefit varying between regions (*high confidence*). {6.6, 6.7, Box TS.7}.

⁴⁶ A general term for how the climate system responds to a radiative forcing (see Glossary).

⁴⁷ The choice of emissions metric depends on the purposes for which gases or forcing agents are being compared. This report contains updated emission metric values and assesses new approaches to aggregating gases.

⁴⁸ For other GHGs, there was insufficient literature available at the time of the assessment to assess detectable changes in their atmospheric growth rate during 2020.

⁴⁹ Near term: (2021–2040)

D.2.3 Scenarios with very low or low GHG emissions (SSP1-1.9 and SSP1-2.6) would have rapid and sustained effects to limit human-caused climate change, compared with scenarios with high or very high GHG emissions (SSP3-7.0 or SSP5-8.5), but early responses of the climate system can be masked by natural variability. For global surface temperature, differences in 20-year trends would *likely* emerge during the near term under a very low GHG emission scenario (SSP1-1.9), relative to a high or very high GHG emission scenario (SSP3-7.0 or SSP5-8.5). The response of many other climate variables would emerge from natural variability at different times later in the 21st century (*high confidence*). {4.6, Cross-Section Box TS.1} (**Figure SPM.8, Figure SPM.10**)

D.2.4 Scenarios with very low and low GHG emissions (SSP1-1.9 and SSP1-2.6) would lead to substantially smaller changes in a range of CIDs³⁶ beyond 2040 than under high and very high GHG emissions scenarios (SSP3-7.0 and SSP5-8.5). By the end of the century, scenarios with very low and low GHG emissions would strongly limit the change of several CIDs, such as the increase in the frequency of extreme sea level events, heavy precipitation and pluvial flooding, and exceedance of dangerous heat thresholds, while limiting the number of regions where such exceedances occur, relative to higher GHG emissions scenarios (*high confidence*). Changes would also be smaller in very low compared to low emissions scenarios, as well as for intermediate (SSP2-4.5) compared to high or very high emissions scenarios (*high confidence*). {9.6, Cross-Chapter Box 11.1, 11.2, 11.3, 11.4, 11.5, 11.6, 11.9, 12.4, 12.5, TS.4.3}



CITY OF CARMEL-BY-THE-SEA

Climate Committee

Staff Report

August 18, 2021
ORDERS OF BUSINESS

TO: Climate Committee Members

SUBMITTED BY: Agnes Martelet, Environmental Compliance Manager

SUBJECT: Update on Climate Action Plan Tasks and Next Steps

RECOMMENDATION:

Receive an update on Climate Action Plan tasks and next steps.

BACKGROUND/SUMMARY:

In April 2021, the City Council adopted a budget amendment that included \$20,000 to hire a consultant to provide technical support to the Climate Committee with greenhouse gas emissions forecasting and reduction strategies. A Request for Proposals was issued and a consultant, LSA Associates, was selected. A Professional Services Agreement with LSA is currently being finalized. The Environmental Compliance Manager will provide an update to the Committee on the consultant selection, proposed tasks, schedule, and next steps.

FISCAL IMPACT:

In April 2021, the City Council adopted a budget amendment that included \$20,000 for greenhouse gas emissions technical support. This funding was carried over to the Fiscal Year 2021-2022 budget.

ATTACHMENTS:



CITY OF CARMEL-BY-THE-SEA

Climate Committee

Staff Report

August 18, 2021
ORDERS OF BUSINESS

TO:	Climate Committee Members
SUBMITTED BY:	Agnes Martelet, Environmental Compliance Manager
SUBJECT:	Review, provide comments, and/or approve the Summary Sheet for the Monterey Regional Waste Management District Presentation

RECOMMENDATION:

Review, provide comments, and/or approve the Summary Sheet for the Monterey Regional Waste Management District Presentation

BACKGROUND/SUMMARY:

The Monterey Regional Waste Management District provided a presentation on their operations and efforts to increase waste diversion from the Marina landfill. The summary for the presentation is included in Attachment 1.

FISCAL IMPACT:

N/A

ATTACHMENTS:

Attachment 1: MRWMD Summary Sheet



CITY OF CARMEL-BY-THE-SEA

Monterey Regional Waste Management District Operations

January 21, 2021 Meeting

TO:	Climate Committee Members
SUBMITTED BY:	Carrie Theis, Committee Member

SUMMARY

Asset:

Regional recycling, waste processing and disposal facility

Hazards:

Methane gas emissions

General Comments and Outlook:

The Monterey Regional Waste Management District is located north of Marina and has been operating since 1951 to handle the waste processing and disposal along with recycling for the Monterey Peninsula. Over the years they have added programs to their facility producing an integrated approach to reduce, reuse and recycle the waste produced on the Monterey Peninsula.

The decomposing of waste materials and organics creates methane gas. Extraction of methane from the landfill creates renewable power for the use by the District and Monterey 1 Water (M1W) to help produce recycled water. Soon the District will be converting landfill gas to Compressed Natural Gas (CNG) to provide fuel for the trash trucks. CNG is a carbon negative fuel. The extraction of methane gas from the landfill produces 5 megawatts of power and the CO2 offset is about 22,000 tons per year.

Windrow Composting is used to manage organics processing to produce compost for nearby organic farms. This process keeps the yard trimmings and organics out of the landfill and creates a nutrient rich compost. It has been shown that applying ¼" of compost (California Carbon Project) to grasslands will improve the soil environment for over a 100 years. Plants thrive more by pulling CO2 out of the air to help offset greenhouse gas emissions. A more widespread application of compost could work to reverse climate change by becoming a carbon bank.

Food recovery and organics processing has become a priority for the State of California with the implementation of AB1826 in 2014 for the mandatory commercial organics recycling and SB1383 in 2016 to mitigate climate change with the Short-Lived Climate Pollutant Reduction Strategy.

The goal of SB1383 is to divert 75% of organic material from the landfill and 20% of edible food for recovery by 2025. To try to obtain the goal of 75% diversion of organic food waste, the waste haulers will be implementing the collection of residential food waste sometime this year. To handle the increased tonnage of organic materials, the Waste Management District will need to implement a larger anaerobic digesting system to prevent the methane gas from being released into the environment. The edible food recovery part of this bill will be accomplished by having the restaurants partner with the Monterey County Food Bank.

Identified Issues:

Attachment 1

- The challenges will be to meet the diversion goals set by the State by continuing to educate the public about recycling and introduce the concept of organics waste diversion. The program logistics of diverting edible food from the restaurants to the Monterey Food Bank program is yet to be determined.

REFERENCES

- Monterey Regional Waste Management District website – www.mrwmd.org.
- Monterey Regional Waste Management District presentation to the Climate Committee: <https://www.youtube.com/channel/UCAu0eILwM1JCDkzD7Js86mA>

History

- Version 1 presented at committee meeting on 8/19/21



CITY OF CARMEL-BY-THE-SEA

Climate Committee

Staff Report

August 18, 2021
ORDERS OF BUSINESS

TO: Climate Committee Members

SUBMITTED BY: Agnes Martelet, Environmental Compliance Manager

SUBJECT: Review and Update the List of Partners to Engage and Discuss the Public Outreach Approach for the Climate Action and Adaptation Strategy Development Phase

RECOMMENDATION:

Review and Update the List of Partners to Engage and Discuss the Public Outreach Approach for the Climate Action and Adaptation Strategy Development Phase

BACKGROUND/SUMMARY:

The Climate Committee previously assembled a list of community organizations and regional partners to engage, including outreach leads for each organization. As the project is moving into the climate adaptation and action strategy phase, the Committee should reevaluate and update the outreach list to determine stakeholders that should be involved in this phase. A draft list is provided in Attachment 1.

Climate Committee members should also discuss how best to reach out to these stakeholders to get meaningful feedback on potential strategies.

FISCAL IMPACT:

N/A

ATTACHMENTS:

Attachment 1: Community Organizations and Regional Partners

Community Organizations and Regional Partners to Engage
City of Carmel-by-the-Sea Climate Project

August 2021

	ORGANIZATION	Topic (s)	Contact Persons	Committee Lead	Previously Contacted?	Contact for Strategy Outreach?
Utilities						
1*	Carmel Area Wastewater District	Sewer system Wastewater Treatment Facility	Barbara Buikema, Daryl Lauer	Jeff Baron, Agnes Martelet	Y	
2	Central Coast Community Energy (was Monterey Bay Community Power)	Energy supply	Dan Bertoldi J.R. Killigrew	Evan Kort, Agnes Martelet	Y	
3	Cal Am	Water supply		Michael LePage		
4	Monterey Peninsula Water Management District	Water supply	Stephanie Locke	Michael LePage	Y	
5	GreenWaste Recovery	Waste management and recycling	Jim Moresco	Carrie Theis, Agnes Martelet		
6	Monterey Regional Waste Management District	Waste management and recycling	Tim Flanagan	Carrie Theis	Y	
7	PG&E	Energy supply	Jeana Arnold Teri Vetere	Carrie Theis, Jeff Baron	Y	
Professional Organizations						
1	California AIA	Built environment resilience	Libby Barnes	John Hill	Y	X
2	Chamber of Commerce	Local business, tourism	Jenny McMurdo	Carrie Theis	Y	X
3	Monterey County Association of Realtors	Sea Level Rise Wildfire risk	Ben Beasley	Scott Lonergan, LaNette Zimmerman	Y	X
4	MCCVB	Visitors		Carrie Theis		
5	Visit Carmel	Visitors / local business	Amy Herzog	Carrie Theis	Y	X
6	Sunset Center	Resilience	Christine Sandin	LaNette Zimmerman		

	ORGANIZATION	Topic (s)	Contact Persons	Committee Lead	Previously Contacted?	Contact for Strategy Outreach?
Local Community Groups						
1	Carmel Residents Association	Community resilience Climate action	Fred Bologna Nancy Twomey	Jeff Baron	Y	X
2	Carmel Rotary Club	Community resilience Climate action	Susan Prest	Carrie Theis, Jeff Baron	Y	X
3	Carmel Lions Club	Community resilience Climate action	Heidy Mozingo	Carrie Theis		X
4	Friends of Carmel Forest	Urban forest	Ramie Allard	Scott Lonergan		X
5	Friends of MTNP	Wildfire risk Sensitive habitats	Karen Ferlito Greg D'Ambrosio	Scott Lonergan	Y	X
6	Del Monte Forest Conservancy	Wildfire risk		Scott Lonergan	Y	
7	Landwatch	Transportation, housing	Mike DeLapa	Jeff Baron	Y	X
8	Sustainable Carmel	Climate action	Ellen Gannon	Agnes Martelet		X
9	Church auxiliaries	Community resilience		LaNette Zimmerman	Y	X
10	Heritage Society	Community resilience	Karl Iverson	Michael LePage		X
11	Forest Theater Guild	Community resilience	Steven Moore	LaNette Zimmerman Carrie Theis		
12	PacRep	Community resilience	Steven Moore	LaNette Zimmerman Carrie Theis		
13	Carmel Women's Club	Community resilience Climate action	Nancy Twomey	Jeff Baron		X
City Departments						
1	Police	Emergency preparedness Transportation	Paul Tomasi	Agnes Martelet	Y	X
2	Fire	Emergency preparedness	Carmyn Prew, Sichel Young	John Hill	Y	X
3	Forestry Division (PW)	Urban forest Sensitive habitats	Sara Davis	Agnes Martelet	Y	X
4	Public Works	Storm Drain Master Plan	Bob Harary	Agnes Martelet	Y	X
5	Library	Historic events	Katie O'Connell	Evan Kort	Y	

	ORGANIZATION	Topic (s)	Contact Persons	Committee Lead	Contacted?	Contact for Strategy Outreach?
City Commissions						
1	Forest & Beach Commission	Sea Level Rise Urban forest Sensitive habitats Climate action		Scott Lonergan	Y	X
2	Planning Commission	Built environment resilience Climate action		Michael LePage		X
3	City Traffic Safety Committee	Transportation		Agnes Martelet	Y	
Federal / State / Regional Agencies						
1	AMBAG	Climate Action Transportation	Amaury Berteaud	Agnes Martelet	Y	X
2	USGS	Sea Level Rise	Patrick Barnard	Agnes Martelet	Y	
3	CalFire	Wildfire risk and resilience Urban forest management	John Reynolds	John Hill	Y	
4	CA Department of Insurance	Wildfire risk / home insurability	Ricardo Lara	John Hill	Y	
5	Fire Safe Council for Monterey County	Michael Emmett		John Hill	Y	
6	California Coastal Commission	Sea Level Rise		Jeff Baron, Agnes Martelet		X
7	IRWMP Group	Water supply Watershed / storm water projects		Michael LePage		
8	Monterey County (OES)	Resilience planning	Kelsey Scanlon	Agnes Martelet	Y	X
9	Monterey County	Sustainability Climate Action	Ashley Paulsworth	Agnes Martelet	Y	X
10	Monterey Bay National Marine Sanctuary (NOAA)	Marine Sanctuary impacts Carmel Beach	Karen Grimmer	Agnes Martelet	Y	X
11	Transportation Agency of Monterey County	Transportation		Jeff Baron		
12	MST	Transportation		Jeff Baron		X
13	US Navy / Coast Guard	Coastal impacts		Carrie Theis		

	ORGANIZATION	Topic (s)	Contact Persons	Committee Lead	Contacted?	Contact for Strategy Outreach?
Educational Groups						
1	Carmel High School Environmental Club	Internships Climate Action partnership / Green Schools	Ellen Gannon	LaNette Zimmerman Scott Lonergan	Y	
2	Stevenson, York, Santa Catalina	Climate Action partnership / Green Schools		LaNette Zimmerman Scott Lonergan	Y	
3	Youth Center	Climate Action partnership / Green Schools		LaNette Zimmerman, Scott Lonergan		
4	CSUMB	Internships Climate Action partnership / Green Schools		LaNette Zimmerman, Scott Lonergan	Y	
5	MPCC	Internships Climate Action partnership / Green Schools		LaNette Zimmerman, Scott Lonergan		
6	MIISS	Climate Action partnership / Green Schools		LaNette Zimmerman, Scott Lonergan	Y	
7	NPS	Climate Action partnership / Green Schools	Ann E. Rondeau (ret.) (pao@nps.edu)	LaNette Zimmerman, Scott Lonergan		
Regional Non-Profit Organizations						
1	Monterey Bay Aquarium	Marine Sanctuary impacts Climate Action Support		Carrie Theis		
2	Ecology Action	Climate Action Support	Kirsten Liske	Agnes Martelet		
Other Stakeholders						
1	Pebble Beach Company	Bluffs / Seawalls / Beach Regional transportation Emergency response		Jeff Baron, Carrie Theis		
2	Principals Involved in 1983 response; authors of the CBTS Shoreline Management Plan	Bluffs / Seawalls / Beach	Greg D'Ambrosio (past Assistant City Administrator); David Shonman (Coastal Biologist)	Evan Kort, Scott Lonergan	Y	
3	Cities with similar challenges (e.g. PG, Monterey, Seaside, Pacifica, Del Mar, Malibu)			Jeff Baron, Carrie Theis		